

Behavior of Some 2(3*H*)-Furanones Bearing A Chromone Moiety as Alkylating Agents

A.K.El-Ziaty, W.S.I.Abou-Elmagd, S.K. Ramadan* and A.I. Hashem

Chemistry Department, Faculty of Science, Ain Shams University, Abassia, Cairo, Egypt, 11566.

4-OXO-4*H*-chromen-3-carboxaldehyde (2) condensed with 3-arylpropionic acids (1a-c) in the presence of thionyl chloride/*N,N*-dimethylformamide mixture as a cyclodehydrating agent to yield 5-aryl-3-chromonylmethylene-2(3*H*)-furanones 3a-c as mixtures of (*E*) and (*Z*) stereoisomers. These furanones treated with the Lewis acid AlCl₃ in benzene, toluene and chlorobenzene to furnish 4,4-diaryl-1-(3-chromonyl)buta-1,3-diene-2-carboxylic acids 4a-f as mixtures of geometrical (*E,E*- and *E,Z*-) stereoisomers via an intermolecular alkylation mode. The unfavored intramolecular alkylation is explained on the basis of the decreased electron density at C₂ of the chromone moiety causing the attack of the intermediate carbocation on this position becomes difficult.

Keywords: 2(3*H*)-Furanones, Chromone-3-carboxaldehyde, Intermolecular alkylation, Intramolecular alkylation and Butadienecarboxylic acids.


2(3*H*)-Furanones represent a group of heterocycles of special interest. This is due to the facile opening of the lactone ring by both nucleophilic and electrophilic reagents. During the last decades, our research group were interested in the conversion of these furanones into a variety of heterocyclic systems of synthetic and biological importance, *e.g.* pyrrolones, pyrazoles, pyridazinones, triazoles, oxadiazoles, thiazoles and isothiazoles⁽¹⁻²⁸⁾. Also, the behavior of these compounds as alkylating agents has been extensively studied by us. It was found that 2(3*H*)-furanones having an aryl or heteryl group attached to an exocyclic double bond at position-3, when treated with AlCl₃ as the Lewis acid in benzene, toluene and anisole, undergo an alkyl-oxygen bond cleavage to give resonance-stabilized carbocations. The carbocation formed in each case either attacks the *o*-position of the aryl or heteryl group situated at position-3 intramolecularly, or is attacked by the solvent in an intermolecular fashion to produce the corresponding butadienecarboxylic acid. It was possible, via the intramolecular alkylation mode, to obtain fluoranthene⁽²⁴⁾, benzofuran⁽²⁵⁾, benzothiophene⁽²⁷⁾, carbazole⁽²⁹⁾ and quinoline⁽³⁰⁾ derivatives.

*E-mail: sayed_karam88@yahoo.com

From our previous studies, it could be concluded that the fate of the carbocation intermediate, *i.e.* its intra- or inter-molecular alkylation depends mainly on three factors, (i) The nature of the substituents situated at position-3, In other words, the availability of electron density at the *o*-position of this substituent. Definitely higher electron density favors the intramolecular mode. (ii) the nucleophilicity of the solvent, for instance it was found that changing the solvent from benzene to the more nucleophilic solvents toluene and anisole changed the reaction pathway from intramolecular to intermolecular mode⁽²⁶⁾. (iii) the stability of the carbocation intermediate, it was found that a more stable carbocation, being more selective favors the formation of cyclic products via intramolecular alkylation⁽²⁹⁾.


As continuation of our previous studies on the chemistry of furanones, we wish to report herein on the study of behavior of 2(3*H*)-furanones bearing a chromonyl group as alkylating agents. Thus, condensation of 4-oxo-4*H*-chromen-3-carboxaldehyde⁽³¹⁾ with 3-arylpropionic acids (1a-c) using thionyl chloride in *N,N*-dimethylformamide as a cyclodehydrating agent⁽³²⁾ was found to afford the corresponding furanones, 5-aryl-3-chromonylmethylene-2(3*H*)-furanones (3a-c). The structures of these products were inferred from their analytical and spectral data. The IR spectra of compounds(3a-c) revealed the characteristic absorption bands for the lactone C=O at 1763-1756 cm⁻¹ as well as the chromone C=O at 1652-1648 cm⁻¹(*Cf.* the Experimental part).

When the furanones 3a-c were treated with AlCl₃ Lewis acid in the presence of benzene, toluene, chlorobenzene, ring opening occurred with the formation of the corresponding butadienecarboxylic acids (4a-f)(Scheme 1).



Scheme 1.

The structures of (4a-f) were determined on the basis of their analytical and spectroscopic data. The IR spectra of compounds (4a-f) were devoid from the absorption of the lactone C=O group and displayed the characteristic broad absorption band of O-H at 3430-3400 cm^{-1} as well as C=O group for the carboxylic group at 1700-1681 cm^{-1} and chromone C=O at 1667-1630 cm^{-1} . The $^1\text{H-NMR}$ spectra of compounds (4b,c,f) showed duplicate signals of different integrations for H_b proton and protons of CH_3 group for compounds (4b,f) which represent a good evidence for their existence as mixtures of two geometrical *E,E*- and *E,Z*-stereoisomers. The higher δ value for proton H_a as compared with H_b , in $^1\text{H-NMR}$ spectra, was attributed to that H_a is located in β -position of α,β -unsaturated system which make it downfield. The mass spectra exhibited the correct molecular ion peak beside some of abundant peaks (Cf. the Experimental part). Evidently, the butadienecarboxylic acids (4a-f) are formed via the intermediacy of the resonance-stabilized carbocation [A] formed by alkyl-oxygen cleavage of the furanone nucleus. The reaction is then completed by attack of the solvent: benzene, toluene or chlorobenzene in an intermolecular fashion to furnish the acids (4a-f).



[A]

It is to be mentioned that when the reaction was carried out in tetrachloroethane or nitrobenzene as solvents, no products of intramolecular alkylation were isolated. We believe that this unfavored intramolecular mode of reaction is not unexpected since C_2^* in the chromone moiety is a part of α,β -unsaturated system. This results in a decrease of the electron density at this position to an extent that attack of the carbocationic center on this position becomes difficult.

Assignment of the configuration *cis*- (H_b/Ar) for the acids **4** is based on that the most stable configuration of the intermediate carbocation [A] is in *E*-configuration. In such configuration, the extended conjugation increases the stability of this carbocation. The attacking group: phenyl, *p*-tolyl or *p*-chlorophenyl should therefore be *trans*- to the H_b -atom.

Experimental

Melting points were measured on a Gallen Kamp electric melting point apparatus. The IR spectra were recorded using potassium bromide disks on a

Fourier transform infrared Thermo Electron Nicolet 7600 spectrometer (Thermo Fisher Scientific Inc., Waltham, MA, USA) at the Central Laboratory of Faculty of Science, Ain Shams University. The $^1\text{H-NMR}$ spectra were run at 400 MHz on a GEMINI 400 BB NMR spectrometer (GEMINI, Manufacturing & Engineering Inc., Anaheim, CA, USA) using tetramethylsilane (TMS) as internal standard in deuterated dimethylsulphoxide ($\text{DMSO-}d_6$) at the Main Defense Chemical Laboratory, Cairo. The mass spectra were recorded on a Shimadzu GC-MS-QP-1000EX mass spectrometer (Shimadzu Scientific Instruments, Inc., USA) operating at 70 eV at the Micro analytical Center of Cairo University. The reactions were monitored by the thin layer chromatography using Merck Kiesel gel 60F₂₅₄ analytical sheets obtained from Fluka.

General procedure for the synthesis of 5-aryl-3-(4-oxo-4H-chromen-3-yl)methylenefuran-2(3H)-ones 3a-c

(i) *(Chlorosulfonyloxy)-N,N-dimethylmethaniminium chloride (the cyclodehydrating agent)*

Into a 25-ml dropping funnel containing benzene (5 ml) and *N,N*-dimethylformamide (1 ml, 10.2 mmol), thionyl chloride (0.8 ml, 11 mmol) was added. After 5 min, the two phases were separated, and the reagent (lower layer) was used in the next step.

(ii) To a stirred solution of 3-arylpropionic acid (1a-c) (10 mmol) in dichloromethane (25 ml) at 0°C, the cyclodehydrating agent (10 mmol) prepared in step (i) was added. Stirring was continued for 15 min. 4-oxo-4H-chromen-3-carboxaldehyde (10 mmol) was added followed by triethylamine (30 mmol) in dichloromethane (15 ml). The resulting mixture was stirred at room temperature for 5 hr. The organic layer was washed with water (2 × 50 ml) and dried over anhydrous sodium sulfate and then evaporated to give 2(3H)-furanones (3a-c).

3-((4-Oxo-4H-chromen-3-yl)methylene)-5-phenylfuran-2(3H)-ones 3a

Yellow crystals; mp. 233-235°C (benzene). Yield 70%. Anal. Calcd. for $\text{C}_{20}\text{H}_{12}\text{O}_4$ (316.07): C, 75.94; H, 3.82. Found: C, 75.82; H, 3.71. IR (KBr) (ν , cm^{-1}): 3115, 3070 (aryl-H), 1763 ($\text{C}=\text{O}_{\text{Furanone}}$), 1652 ($\text{C}=\text{O}_{\text{Chromone}}$), 1610 ($\text{C}=\text{C}$). $^1\text{H-NMR}$ ($\text{DMSO-}d_6$): δ (ppm) (*E*-form, 75%) 9.09 (s, 1H, $\text{C-H}_{\text{Chromone}}$), 8.18 (s, 1H, H_a), 7.88-7.77 (m, 4H, Ar- $\text{H}_{\text{Chromone}}$), 7.74-7.39 (m, 6H, $5\text{H}_{\text{Phenyl}}$ + $1\text{H}_{\text{Furanone}}$); (*Z*-form, 25%) δ 8.16 (s, 1H, H_a). MS, m/z (%): 316 (M^+ , 6), 288 (7), 183 (3), 155 (4), 111 (12), 105 (81), 92 (24), 77 (100).

3-((4-Oxo-4H-chromen-3-yl)methylene)-5-(4-tolyl)furan-2(3H)-ones 3b

Yellow crystals; mp. 264-266°C (benzene). Yield 79%. Anal. Calcd. for $\text{C}_{21}\text{H}_{14}\text{O}_4$ (330.09): C, 76.35; H, 4.27. Found: C, 76.24; H, 4.11. IR (KBr) (ν , cm^{-1}): 3126, 3028 (aryl-H), 2914 (alkyl-H), 1761 ($\text{C}=\text{O}_{\text{Furanone}}$), 1650 ($\text{C}=\text{O}_{\text{Chromone}}$), 1611 ($\text{C}=\text{C}$). $^1\text{H-NMR}$ ($\text{DMSO-}d_6$): (*E*-form, 70%) δ (ppm) 9.07

(s, 1H, C-H_{Chromone}), 8.18 (s, 1H, H_a), 7.91-7.72 (m, 4H, Ar-H_{Chromone}), 7.59-7.33 (m, 5H, 4H_{Tolyl} + 1H_{Furanone}), 2.38 (s, 3H, CH₃); (Z-form, 30%) δ 8.16 (s, 1H, H_a). MS, *m/z* (%): 330 (M⁺, 2), 292 (13), 276 (10), 237 (8), 194 (9), 177 (26), 149 (14), 120 (20), 119 (100), 105 (37), 91 (21), 77 (64).

5-(4-Chlorophenyl)-3-((4-oxo-4H-chromen-3-yl)methylene)furan-2(3H)-ones 3c

Yellow crystals; mp.308-310°C (benzene).Yield 74%.Anal.Calcd.for C₂₀H₁₁O₄Cl (350.03): C, 68.49; H, 3.16, Cl, 10.11. Found: C, 68.35; H, 3.04, Cl, 10.01. IR (KBr) (ν , cm⁻¹): 3127, 3084 (aryl-H), 1756 (C=O_{Furanone}), 1648 (C=O_{Chromone}), 1613 (C=C). ¹H-NMR (DMSO-*d*₆): (*E*-form, 75%) δ (ppm) 9.05 (s, 1H, C-H_{Chromone}), 8.16 (s, 1H, H_a), 7.89-7.72 (m, 4H, Ar-H_{Chromone}), 7.59-7.39 (m, 5H, 4H_{Chlorophenyl} + 1H_{Furanone}); (Z-form, 25%) δ 8.16 (s, 1H, H_a). MS, *m/z* (%): 352 (M+2, 8), 350 (M⁺, 25), 324 (6), 322 (17), 296 (2), 294 (5), 211 (7), 173 (8), 141 (30), 139 (100), 111 (58), 92 (25), 75 (27).

General procedure for the preparation of 2-((4-oxo-4H-chromen-3-yl)methylene)-4,4-diarylbut-3-enoic acid 4a-f

To a stirred solution of anhydrous aluminum chloride (0.03 mol) in dry benzene, toluene or chlorobenzene (100 ml), a solution of furanones (3a-c) (0.01 mol) in dry benzene, toluene or chlorobenzene was added dropwise at 10-20°C. After complete addition, the whole mixture was stirred at room temperature for further 6 hr, heated on water bath for 2 hr, then left to stand overnight. The complex formed was decomposed with ice cold hydrochloric acid (50 ml) and hence steam-distilled to remove the excess of organic solvent. The residue was triturated with ether, filtered off, and recrystallized from the suitable solvent.

2-((4-Oxo-4H-chromen-3-yl)methylene)-4,4-diphenylbut-3-enoic acid 4a

White crystals; mp.258-260°C(Petroleum ether, 60-80°C/benzene, 1:1).Yield 31%.Anal.Calcd.for C₂₆H₁₈O₄ (394.12):C, 79.17; H, 4.60. Found: C, 79.09; H, 4.51. IR (KBr) (ν , cm⁻¹): 3420 (OH), 3068 (aryl-H), 1700 (C=O_{acid}), 1630 (C=O_{Chromone}), 1610 (C=C). ¹H-NMR (DMSO-*d*₆): δ (ppm) 12.63 (s, 1H, COOH, *D*₂O-exchangeable), 8.04 (s, 1H, C-H_{Chromone}), 7.86-7.47 (m, 4H, Ar-H_{Chromone}), 7.36 (s, 1H, H_a), 7.28-7.04 (m, 10H, 2 Ar-H_{phenyl}), 7.02 (s, 1H, H_b). MS, *m/z* (%): 394 (M⁺, 3), 350 (12), 322 (5), 293 (4), 237 (9), 197 (13), 151 (12), 139 (38), 111 (100), 92 (53), 76 (46), 63 (58).

2-((4-Oxo-4H-chromen-3-yl)methylene)-4-phenyl-4-p-tolylbut-3-enoic acid 4b

Faint brown crystals; mp.128-130°C(benzene).Yield 35%.Anal.Calcd.for C₂₇H₂₀O₄ (408.14): C, 79.40; H, 4.94. Found: C, 79.31; H, 4.85. IR (KBr) (ν , cm⁻¹): 3426 (OH), 3078 (aryl-H), 1682 (C=O_{acid}), 1635 (C=O_{Chromone}), 1615 (C=C). ¹H-NMR (DMSO-*d*₆): (*E,Z*-form, 60%) δ (ppm) 12.70 (s, 1H, COOH, *D*₂O-exchangeable), 8.14 (s, 1H, C-H_{Chromone}), 8.08-7.50 (m, 4H, Ar-H_{Chromone}), 7.37 (s, 1H, H_a), 7.17-7.06 (m, 5H, Ar-H_{phenyl}), 7.00-6.91 (m, 4H, Ar-H_{Tolyl}), 6.89 (s, 1H, H_b), 2.21 (s, 3H, CH₃); (*E,E*-form, 40%) δ 6.87 (s, 1H, H_b), 2.17 (s, 3H,

CH₃).MS, *m/z* (%): 408 (M⁺, 1), 393 (2), 334 (46), 289 (31), 229 (70), 183 (34), 159 (28), 105 (100), 92 (22), 65 (21).

4-(4-Chlorophenyl)-2-((4-oxo-4H-chromen-3-yl)methylene)-4-phenylbut-3-enoic acid 4c

Yellow crystals; mp.208-209°C(Petroleum ether, 60-80°C/benzene, 1:1).Yield 28%.Anal.Calcd.for C₂₆H₁₇ClO₄ (428.08): C, 72.82; H, 4.00; Cl, 8.27. Found: C, 72.73; H, 3.89; Cl, 8.19. IR (KBr) (ν , cm⁻¹): 3430 (OH), 3078 (aryl-H), 1701 (C=O_{acid}), 1652 (C=O_{Chromone}),1619 (C=C). ¹H-NMR (DMSO-*d*₆):(*E,Z*-form, 65%) δ (ppm) 12.63 (s, 1H, COOH, *D*₂*O*-exchangeable), 8.52 (s, 1H,C-H_{Chromone}), 8.24-7.75 (m, 4H, Ar-H_{Chromone}), 7.64-7.39 (m, 4H, Ar-H_{Chlorophenyl}), 7.35 (s, 1H,H_a), 7.17-6.85 (m, 5H, Ar-H_{Phenyl}),6.83 (s, 1H, H_b);(*E,E*-form, 35%) δ 6.65 (s, 1H, H_b).MS, *m/z* (%): 430 (M+2, 4), 428 (M⁺, 11), 411 (13), 409 (12), 383(44), 381 (43), 347 (10), 289 (11), 262 (25), 237 (61), 235 (91), 202 (25), 199 (130), 165 (100), 146 (54), 121 (71), 92 (34), 77 (39).

2-((4-Oxo-4H-chromen-3-yl)methylene)-4,4-dip-tolylbut-3-enoic acid 4d

Faint brown crystals; mp.136-138°C (Petroleum ether, 60-80°C/benzene, 1:1).Yield 38%.Anal.Calcd.for C₂₈H₂₂O₄ (422.15): C, 79.60; H, 5.25. Found: C, 79.57; H, 5.22. IR (KBr) (ν , cm⁻¹): 3425 (OH), 3077 (aryl-H), 1682 (C=O_{acid}), 1636 (C=O_{Chromone}), 1615 (C=C). ¹H-NMR (DMSO-*d*₆): δ (ppm) 12.69 (s, 1H, COOH, *D*₂*O*-exchangeable), 8.13 (s, 1H,C-H_{Chromone}), 8.08-7.48 (m, 4H, Ar-H_{Chromone}), 7.38 (s, 1H,H_a), 7.22-6.89 (m, 8H, 2 Ar-H_{Tolyl}), 6.86 (s, 1H, H_b), 2.20 (s, 6H, 2 CH₃).MS, *m/z* (%): 422 (M⁺, 2), 367 (3), 334 (44), 288 (13), 228 (38), 184 (19), 160 (12), 105 (100), 91 (28), 65 (20).

4,4-Bis(4-chlorophenyl)-2-((4-oxo-4H-chromen-3-yl)methylene)but-3-enoic acid 4e

Yellow crystals; mp.235-237°C(benzene).Yield 25%.Anal.Calcd.for C₂₆H₁₆Cl₂O₄ (462.04): C, 67.40; H, 3.48; Cl, 15.30. Found: C, 67.31; H, 3.38; Cl, 15.22. IR (KBr) (ν , cm⁻¹): 3400-3200 (OH), 3063 (aryl-H), 1695 (C=O_{acid}), 1667 (C=O_{Chromone}), 1625 (C=C). ¹H-NMR (DMSO-*d*₆): δ (ppm) 10.33 (s, 1H, COOH, *D*₂*O*-exchangeable), 8.02 (s, 1H,C-H_{Chromone}), 7.62-7.35 (m, 4H, Ar-H_{Chromone}), 7.50 (s, 1H,H_a),7.23-6.76 (m, 8H,2 Ar-H_{Chlorophenyl}),6.73 (s, 1H, H_b).MS, *m/z* (%): 462 (M⁺, 1), 323 (2), 288 (5), 238 (12), 173 (17), 146 (18), 123 (56), 105 (100), 92 (36), 77 (29).

4-(4-Chlorophenyl)-2-((4-oxo-4H-chromen-3-yl)methylene)-4-p-tolylbut-3-enoic acid 4f

Faint brown crystals; mp.140-142°C(Petroleum ether, 60-80°C/benzene, 1:1).Yield 34%.Anal.Calcd.for C₂₇H₁₉ClO₄ (442.10): C, 73.22; H, 4.32; Cl, 8.00. Found: C, 73.11; H, 4.22; Cl, 7.87. IR (KBr) (ν , cm⁻¹): 3410 (OH), 3078 (aryl-H), 1681 (C=O_{acid}), 1636 (C=O_{Chromone}), 1615 (C=C). ¹H-NMR (DMSO-*d*₆): (*E,Z*-form, 75%) δ (ppm) 12.68 (s, 1H, COOH, *D*₂*O*-exchangeable), 8.15 (s, 1H,C-H_{Chromone}), 8.13-7.79 (m, 4H, Ar-H_{Chromone}), 7.68-7.48 (m, 4H, Ar-

$H_{\text{Chlorophenyl}}$), 7.37 (s, 1H, H_a), 7.16-6.98 (m, 4H, Ar- H_{Tolyl}), 6.93 (s, 1H, H_b), 2.20 (s, 3H, CH_3); (*E,E*-form, 25%) δ 6.90 (s, 1H, H_b), 2.15 (s, 3H, CH_3). MS, m/z (%): 442 (M^+ , missed), 407 (M-Cl, 1), 393 (3), 334 (54), 316 (15), 289 (32), 183 (35), 121 (51), 105 (100), 77 (37), 65 (23).

Refernces

1. **Abou-Elmagd, W. S. I. and Hashem, A. I.**, Synthesis and antitumor activity evaluation of some novel fused and spiro heterocycles derived from a 2(3H)-furanone derivative. *J. Heterocycl. Chem.* **53**, 202-208 (2016).
2. **Abou-Elmagd, W. S. I., El-Ziaty, A. K. and Abdalha, A. A.**, Ring transformation and antimicrobial activity of indolyl-substituted 2(3H)-furanones. *Heterocycl. Commun.* **21**(3), 179-184 (2015).
3. **Hashem, A. I., Abou-Elmagd, W. S. I. and Abd-Elaziz A.**, Synthesis and reactions of some 2(3H)- and 2(5H)-furanone derivatives: a comparative study. *Eur. Chem. Bull.*, **3**(11), 1064-1068 (2014).
4. **Abou-Elmagd, W. S. I. and Hashem, A. I.**, Conversion of some 2(3H)-furanones into pyrrolinotriazine and oxazolopyrimidine derivatives. *J. Heterocycl. Chem.* **49**, 947-950 (2012).
5. **Abou-Elmagd, W. S. I. and Hashem, A. I.**, Novel synthesis of some isatinhydrazones and pyridazinophthalazines. *Synth. Commun.* **43**, 1083-1088 (2013).
6. **Abou-Elmagd, W. S. I. and Hashem, A. I.**, Synthesis of 1-amidoalkyl-2-naphthols and oxazine derivatives with study of their antibacterial and antiviral activities. *Med. Chem. Res.* **22**, 2005-2010 (2013).
7. **Sayed, H. H., Hashem, A. I., Youssef, N. M. and El-Sayed, W. A.**, Conversion of 3-arylo-5-phenyl-2(3H)-furanone into other heterocycles of anticipated biological activity. *Arch. Pharmazie*, **340**, 315-319 (2007).
8. **Hashem, A. I., Youssef, A. S. A., Kandeel, K. A. and Abou-Elmagd, W. S. I.**, Conversion of some 2(3H)-furanones bearing a pyrazolyl group into other heterocyclic systems with a study of their antiviral activity. *Eur. J. Med. Chem.*, **42**, 934-939 (2007).
9. **Kandeel, K. A., Youssef, A. S. A., Abou-Elmagd, W. S. I. and Hashem, A. I.**, Behaviour of 3-benzylamino-5-aryl-2(3H)-furanones towards some nitrogen nucleophiles. *J. Heterocycl. Chem.*, **43**, 957-962 (2006).
10. **Youssef, A. S. A., Kandeel, K. A., Abou-Elmagd, W. S. I. and Hashem, A. I.**, Conversion of 5-aryl-3-phenylthio-2(3H)-furanones into some nitrogen and sulfur-containing heterocycles. *Phosphorus Sulfur and Silicon*, **182**(1), 85-97 (2006).
11. **El-Kousy, S. M., Hashem, A. I., El-Torgoman, A. M. and Salama G. M.**, Reactions of some arylazofuranones with aminoacids and malononitrile. *Afindad IX*, **503**, 61-64 (2003).

12. **Hamad, A. S., Zaki, M. Y., Eid, N. N. and Hashem, A. I.**, Conversion of 2(3*H*)-furanones into 1,3,4-oxadiazoles. *Heteroatom Chem.*, **14**, 570-574 (2003).
13. **Hamad, A. S. and Hashem, A. I.**, Novel synthesis of *N*-[6-aryl-4-(*E*)-2-furylmethylene-1,2,3,4-tetrahydro-3-oxopyridazin-1-ylcarbonyl]-*p*-toluenesulphonamides and *N*-[5-(*E*)-1-arylmethyl-2-(2-furyl)vinyl]-1,3,4-oxadiazol-2-yl]-*p*-toluenesulphonamides. *J. Heterocycl. Chem.*, **39**, 1325-1328 (2002).
14. **Derbala, H. A., Hamad, A. S., El-Sayed, W. A. and Hashem, A. I.**, Conversion of 3-aryl-5-phenyl-2(3*H*)-furanones into 3(2*H*)-isothiazolone derivatives. *Phosphorus, Sulfur and Silicon*, **175**, 153-162 (2001).
15. **Hamad, A. S., Derbala, H. A., El-Sayed, W. A. and Hashem, A. I.**, Behavior of 2(3*H*)- and 2(5*H*)-furanones having no exocyclic double bond towards some nitrogen nucleophiles. *Acta Chim. Slov.*, **48**, 417-425 (2001).
16. **Hamad, A. S. and Hashem, A. I.**, A facile method for the synthesis of 6-aryl-1-(3-chloropropanoyl)-4-[(*E*)-1-(2-furyl)methylidene]-1,2,3,4-tetrahydropyridazinones and 2-chloroethyl-5-[α -aracyl- β -(2-furyl)-*E*-vinyl]-1,3,4-oxadiazoles. *Molecules*, **5**, 895-907 (2000).
17. **Yassin, S., Abd El-Aleem, A. H., El-Sayed, I. E. and Hashem, A. I.**, Conversion of 1-(α -aracyl- β -(thienyl)acryloylsemicarbazides into 1,2,4-triazole derivatives. *Rev. Roum. Chim.*, **41**, 989-993 (1996).
18. **Yassin, S., Abd El-Aleem, A. H., El-Sayed, I. E. and Hashem, A. I.**, Synthesis of some diaroylhydrazines, 1,3,4-oxadiazoles and pyridazin-3-ones bearing thiophene nucleus. *Collect. Czech. Chem. Commun.* **58**, 1925-1930 (1993).
19. **El-Kousy, S. M., El-Torgoman, A. M., El-Bassiouny, A. A. and Hashem, A. I.**, Synthesis and reactions of some β , γ -butenolides with study of biological activity of the resulting products. *Pharmazie*, **43**(2), 80-81 (1988).
20. **Hashem, A. I. and Shaban, M. E.**, Preparation and reactions of some derivatives of acrylic acid hydrazides and amides. *J. Prakt. Chem.*, **323**, 164-168 (1981).
21. **Hashem, A. I.**, 5-Oxo-2-pyrrolines from β , γ -Butenolides. *J. Prakt. Chem.*, **321**, 516-518 (1979).
22. **Awad, W. I., Hashem, A. I. and El-Badry, K.**, Preparation and reactions of some acyl azides. *Indian J. Chem.*, **13**(B), 1139-1141 (1975).
23. **Hashem, A. I.**, Reactions of Some β , γ -Butenolides. *J. Prakt. Chem.*, **315**, 335-338 (1973).
24. **Hashem, A. I. and Ismail, Z. M.**, Polynuclear compounds from highly conjugated butenolides. *Indian J. Chem.*, **12**(B), 902-904 (1974).
25. **Hashem, A. I.**, Benzofuran Derivatives from β , γ -Butenolides. *J. Prakt. Chem.*, **319**, 689-692 (1977).

26. **Hashem, A. I. and Eid, E. E.**, Behavior of some β,γ -butenolides towards Friedel-Crafts alkylation with benzene, toluene and anisole. *Indian J. Chem.*, **16(B)**, 502-504(1978).
27. **Hashem, A. I., El-Kousy, S. M., El-Torgoman, A. M. and El-Bassiouny, A. M.**, Benzothiophene derivatives from 2(3H)-furanones. *Internat. J. Chem.*, **2(2)**, 87-91 (1991).
28. **Hashem, A. I., Kandeel, K. A., Youssef, A. S. A. and Abou-Elmagd, W. S. I.**, Behavior of some 2(3H)-furanones bearing a pyrazolyl group as alkylating agents. *J. Chem. Res.*, 315-317(2006).
29. **Abou-Elmagd, W. S. I.**, Behaviour of 3-(3-indolylmethylene)-5-aryl-2(3H)-furanones as alkylating agents: intra- versus inter-molecular alkylation. *J. Chem. Res.*, 305-307(2007).
30. **Abou-Elmagd, W. S. I.**, Behaviour of 3-(2-pyridinylmethylene)-5-aryl-2(3H)-furanones and 3-(3-pyridinylmethylene)-5-aryl-2(3H)-furanones as alkylating agents: a comparative study. *J. Heterocycl. Chem.*, **48**, 1361-1365(2011).
31. **Nohara, A., Umetani, T. and Y. Sanno, Y.**, Studies on antianaphylactic agents—I: A facile synthesis of 4-oxo-4H-1-benzopyran-3-carboxaldehydes by Vilsmeier reagents. *Tetrahedron*, **30**, 3553-3561(1974).
32. **Khan, R. H. and Rastogi, R. C.**, A convenient and facile synthesis of 2-arylidene-4-phenylbut-3-en-4-olides by use of *N,N*-dimethyl(chlorosulphonyl)methaninium chloride as cyclodehydrating agent. *J. Chem. Res.*, 260-263(1991).

(Received 3/7/2016;
accepted 18/9/2016)

سلوك بعض مركبات 2(H-3) فيورانونات والحاملة لمجموعة كرومون ككاشف ألكلة

أحمد كامل الزيياتى ، وائل سيد ابراهيم أبوالمجد ، سيد كرم رمضان
وأحمد إسماعيل هاشم
قسم الكيمياء – كلية العلوم – جامعة عين شمس – العباسية – القاهرة –
مصر.

تتفاعل أحماض 3-ارويل بروبيونيك مع 4-اكسوكرومون-3-كربوكسالديهد باستخدام مخلوط كلوريد ثيونيل وثنائي ميثيل فورماميد ككاشف لنزع الماء لتعطى 5-أريل-3-كرومونايل ميثيلين-2(H3)-فيورانونات كمخلوط من نظائر *E*-و-*Z*. وقد تفاعلت هذه الفيورانونات مع حامض لويس (كلوريد الالومنيوم اللامائي) فى وجود البنزين ، الطولوين والكلوروبنزين لتعطى أحماض 4،4-ثنائي أريل-1-(3-كرومونايل)بيوتا-1 و3-دايين-2-كربوكسيليك كمخلوط من نظائر *E*،*E*-و-*Z* عن طريق نظام الألكلة الخارجية. يمكن تفسير نظام الألكلة الداخلية الغير مناسب عن طريق نقص الكثافة الالكترونية عند موضع 2 فى مجموعة الكرومون مما يجعل هجوم الكربوكاتيون صعب عند هذا الموضع.