



Effect of Carboxymethyl Cellulose with Edible Coating on Chemical and Physical Characters of Fresh-Cut Green Bean Pods during Cold Storage.



CrossMark

Huda A. Ibrahim¹, Hayam A. A. Mahdy², Ahmad A. Suliman³ and Mohamed A. A. Abdullah⁴

¹Vegetable Research Dept., National Research Centre, 33 El-Buhouth St., (former El- Tahrir St.,) Dokki, Giza, Egypt. Postal Code: 12622.

²Botany Dept., National Research Center, 33 El-Buhouth St., (former El- Tahrir St.,) Dokki, Giza, Egypt. Postal Code: 12622.

³Horticultural Crops Technology Dept., National Research Centre, 33 El-Buhouth St., (former El- Tahrir St.,) Dokki, Giza, Egypt. Postal Code: 12622.

⁴Vegetable Handling Dept., Horticulture Research Institute, Agricultural Research Center, Giza, Egypt

Abstract

Pods of green beans (*Phaseolus vulgaris* L.) cv. Paulista were harvested and stored in storage. The current study aimed to evaluate the physical characters and chemical constituents of stored pods and treated by carboxymethyl cellulose plus gum arabic 5% for 5 minutes, carboxymethyl cellulose plus calcium chloride 0.5% for 5 minutes, carboxymethyl cellulose plus ascorbic acid 5% for 5 minutes, carboxymethyl cellulose for 5 minutes and tap water for 5 minutes which served as control. Fresh cut green pods wrapped with polypropylene bags and stored under 5°C, 85-90% relative humidity for twenty days, then studying the effect of previous treatments on physical and chemical characters of fresh-cut green bean pods under cold storage conditions. Results indicated that treated pods by Carboxymethyl Cellulose with arabic gum reduced the loss of pod weight, decay score, and maintained pods general appearance, firmness, total soluble solids, total chlorophyll content, and led to a decrease in total phenolic compound and microbial count, followed by with Calcium Chloride, then with Ascorbic Acid up to 20 days of storage compared to untreated control.

Keywords: Green beans, Carboxymethyl cellulose, Gum arabic, Calcium chloride.

1. introduction

Green beans (*Phaseolus vulgaris* L.) are considered one of the most important leguminous crops in Egypt for local consumption and exportation. It is widely used as a protein source and for other nutrients in many developing countries including Egypt (Saleh *et al.*, 2018). In addition, it is rich in vitamins, minerals, and carotene. It is a source of vitamin B, macro-and micro elements (Al-Sanabani *et al.*, 2016), also are climacteric type of respiration pattern; therefore, changes in the flavor, texture and appearance are limiting factors of the quality of pods (Kasim and Kasim, 2015). Generally green beans harvested at a physiologically immature stage. The most rapid of growth at the time of harvest and green beans exhibit comparatively higher respiration, even when held at low temperatures, cutting of green bean also accelerates respiration rate more than in whole beans. Therefore, fresh-cut of green beans highly decrease in quality. In order to, prevent the loss of quality of fresh-cut vegetables, several treatment methods are being practiced reduce the changes that occur after harvesting. Postharvest treatments for fresh produce seek to reduce physiological processes of senescence and maturation, inhibit development of physiological disorders, and minimize the risk of pathogens growth and contamination. In addition to, basic postharvest technology

for temperature management, an array of others has been developed, including various physical (irradiation, heat and edible coatings); chemical (anti-browning, antioxidants and antimicrobials); and gaseous treatments. Edible coating is used to inhibit oxygen, migration of moisture, aromas, carbene dioxide and lipids and to improve the mechanical integrity or handling characteristics of fruits (Röjas-Graü *et al.*, 2011). Carboxymethyl cellulose (CMC) one of the most used and widely common polysaccharides in the food, agriculture, and other industries and has been safety used as a food additive and is approved by the FDA and EFSA (Shebis *et al.*, 2022). Carboxymethyl cellulose prepared by a reaction of cellulose with sodium hydroxide and chloroacetic acids (Biswal and Singh, 2004). In addition, carboxymethyl cellulose can be produced from agricultural waste such as papaya (Rachtanapun *et al.*, 2007), Mimoso pigra (Rachtanapun and Rattanapanone, 2011), durian (Rachtanapun *et al.*, 2011), and pomelo peel. Many research have applied carboxymethyl cellulose for extend shelf-life and storage period of fresh produce such as peach and pear (Tog'rul and Arslan, 2004), avocado (Maftoonazad and Ramaswamy, 2005), mango (Rachtanapun *et al.*, 2008), cucumber (Adetunji *et al.*, 2013), Rambutan (Kasinee *et al.*, 2017) and Strawberry (Shebis *et al.*, 2022). The produced, novel Carboxymethyl

*Corresponding author e-mail: a_elsagheer2006@yahoo.com; (ggggggggggggggggggggg).

Receive Date: 08 November 2023, Revise Date: 14 December 2023, Accept Date: 17 December 2023

DOI: 10.21608/EJCHEM.2023.247222.8831

©2024 National Information and Documentation Center (NIDOC)

cellulose were applied alone and in combination with edible coating, gum arabic, calcium chlorid, and ascorbic acid, known to protect fruits and vegetables. Gum arabic, which is used as an edible coating to preserve the quality of fruits, is a substance that contains a large number of important polysaccharides used in agricultural processing due to its ease of formation in the form of a film, and the unique emulsifying and encapsulating properties that received the highest safety ratings by a joint research between the World Health Organization and the Food Organization Agriculture and Food Additives Committee (Anderson and Eastwood, 1989; Motlagh *et al.*, 2006). When used as an edible coating material, gum arabic also acted on some important consequences of delaying signs of ripeness during storage of apples in storage under cryogenic conditions (El-Anany *et al.*, 2009; Huang *et al.*, 2021). Calcium is as one of important element for application postharvest for several fresh to stop the causes that lead to the occurrence of many factors of post-harvest deterioration and thus to improve the storage conditions of field crops. It also supports many physiological and biological functions within the cell wall, organs, and chromosomes, increasing enzyme activation and Ca-phytohormone interactions. Calcium has other positive effects, most notably in vegetables, so it can extend storage life by reducing respiration rate, delaying signs of ripeness, increasing durability and vitamin C content, and reducing the appearance of storage decay (Naveena and Immanule 2017 and Gameda, 2021). Ascorbic acid is an antioxidant molecule (Shalata and Neumann 2001) which contributes to the detoxification of reactive oxygen species (ROS) and therefore the usage of ascorbic acid is associated with delayed senescence and resistance plants to oxidative stress (Bilgin 2021). This research aimed to study the effects of carboxymethyl cellulose alone or in combination with Edable coating on pod quality of fresh-cut green bean under cold storage conditions.

2. Materials and Methods:

Green bean (*Phaseolus vulgaris* L.) cv. Paulista pods were obtained, from private farm at El Behira governorate during the two successive seasons of 2022 and 2023 respectively. Green pods were harvested at commercial maturity (i.e., bright green, tender fleshy and with small green immature seeds) at the 11th and 16th of January during the two successive seasons. The precooling process was carried out and then green pods were transferred to the postharvest laboratory in Horticulture Research Institute, Giza governorate. Pods without any visual defects were washed with tap water to eliminate soil residues. Pods were cut from both ends, using a sharp stainless steel knife.

The selected pods were of the same length and completely free of damage or visual defects. The pods were distributed into five groups for the following treatments

1. Dipping in the solution of Carboxymethyl Cellulose plus gum Arabic 5% for 5 minutes (CMC plus Gum Arabic).
2. Dipping in the solution of Carboxymethyl Cellulose plus calcium chloride 0.5% for 5 minutes (CMC plus CaCl₂).
3. Dipping in the solution of Carboxymethyl Cellulose plus Ascorbic acid 5% for 5 minutes (CMC plus Ascorbic acid).
4. Dipping in the solution of Carboxymethyl cellulose for 5 minutes (CMC).

5. Dipping in tap water for 5 minutes which served as control.

In the laboratory, Carboxymethyl Cellulose, were prepared by the water hyacinth plants. Washed, dried, and ground. The dried powder was then extracted with 1 % (w/v) sodium hydroxide (NaOH) solution at the ratio of cellulose to solvent being 1: 30 (w/v) at 90 °C for 3 h. The mixture was filtered, washed with water until neutral and dried at 18 °C. The dried cellulose was passed through 100 mesh sieves and kept in the polyethylene bags. Fifteen grams of prepared cellulose was mixed with 450 ml of isopropanol, and 50 ml of 30 % NaOH solution in a beaker for 30 min. Then, 18 g of chloroacetic acid was added into the mixture and stirred for 1.5 h. The mixture was covered with aluminium foil and placed in an oven at 55 °C for 3.5 h. The obtained solid was neutralized with 90 % acetic acid, filtered, washed 5 times with ethanol and 1 time with methanol. The final product CMC was obtained after drying at ambient temperature (Kasinee *et al.*, 2017).

Gum arabic solutions were prepared by dissolving 5 and 10 g of powder gum arabic in 100 mL purified water. The solutions were stirred with low heat (40 °C) for 60 min on a magnetic stirrer/hot plate (Model: HTS-1003), then filtered to remove any undissolved impurities using a vacuum flask. After cooling to 20 °C, glycerol monostearate (1.0%) (Sigma) was added as a plasticizer to improve the strength and flexibility of the coating solutions. The pH of the solutions was adjusted to 5.6 with 1 N NaOH (Ali *et al.*, 2010).

Each treatment consists of 200 g of fresh cut green pods wrapped with polypropylene bags and stored under 5 °C, 85-90% relative humidity. Samples were taken at random from the three replicates for each treatment. The treatments were examined immediately after harvest and every four days intervals for the following parameter and stored for 20 days.

2.1 Weight loss percentage: it was estimated according to the following equation: Weight loss% = [(Initial weight - weight of fruits at sampling date)/Initial weight of fruits] x 100.

2.2. Decay: it was determined as score system of 1= none, 2= slight, 3= moderate, 4= moderately severe, 5= severe. This depends on decay percentage on pods (Watada and Morris, 1996; Jimenez *et al.*, 1998).

2.3. General appearance: it was determined as score system of excellent > 9, good > 7 to 8.9, fair > 5 to 6.9, poor > 3 to 4.9, and unassailable > 2.9. The scale depends on morphological defects such as shriveling, fresh appearance, color change of pods and decay. Pods rating (5) or below considered unmarketable (Watada and Morris, 1996; Jimenez *et al.*, 1998).

2.4. Firmness: the average firmness of the fruits was measured in kg/cm² by digital force Gauge model FGV 50 A, Shimpo Instrument Co, Japan, with total capacity of 20 kg/cm² and resolution of 0.01kg/cm² using cone pointed head.

2.5. Total soluble solids percentage (TSS%) was determined as a composite juice sample by digital refractometer of model Abbe Leica according to (A.O.A.C., 1990).

2.6. Total chlorophyll: was determined as described in (Shehata *et al.*, 2018). In brief, 0.5 g of fresh sample were homogenized with 5 mL dimethyl formamide and kept in the dark in the refrigerator for 48 h. The absorbance was

then measured at 470, 647 and 663 nm with a spectrophotometer (model UV-2401 PC, International Equipment Trading LTD. (IET), Milano, Italia).

2.7. Total phenolic compound: was calculated by using the Folin–Ciocalteu reagent with some alteration by using gallic acid as a standard curve (El-Mogy *et al.*, 2020). Five grams of the sample was diluted using 5 mL of methanol (80%). The solution was blended with 2.5 mL of Folin–Ciocalteu (10-fold with distilled water) and added to 2.5 mL of distilled water. Afterwards, 2 mL of aqueous sodium carbonate solution (7.5%, w/v) was added after incubation for 5 min. The final solution was mixed and incubated in the dark at room temperature for 1 h. The absorption was assessed at 765 nm using the spectrophotometer, and the results were expressed as milligrams of gallic acid equivalent (GAE) per 100 mg of fresh fruit weight.

2.8. Microbiological analysis: each sample was prepared by homogenizing 10 gm of sliced green bean with 100 ml sterile 0.1 peptone water for 2 min. Dilution by 0.1 pepton water was made as needed in Plate Count Agar (PCA) and incubated for 24 hr at 37 °C for the determination of mesophilic aerobic microorganisms. Viable counts were determined by counting the number of colonies and reported as colony forming units per gram, CFU/g (Anese and Nicoli, 1997).

All data were subjected to statistical analysis according to the procedures reported by Snedecor and Cochran (1982) and means were compared by Duncan's multiple range tests (1955).

Results and Discussions

Weight loss percentage

Data in Table (1) indicate increase in weight loss till the end of storage period, where the maximum loss was occurred. The results showed that the weight loss of green bean pods increased significantly and steadily as the storage period increased in the pods collected during the two growing seasons. This phenomenon may be explained by the loss of moisture inside the pods through transpiration and the loss of part of the dry matter content as a result of tissue respiration during storage. These results are consistent with (Zhu *et al.*, 2008, Abdullah *et al.*, 2018a and El-Mogy *et al.*, 2020).

With regard to the effect coefficients, the data showed that there were significant differences between the coefficients in the percentage of weight loss during the storage period. All treatments maintained their normal weight during the storage period when compared to untreated plants. Moreover, dipping green bean pods treated for Carboxymethyl Cellulose with gum arabic 5% for 5 minutes was the most effective treatment in reducing the percentage of weight loss % followed by Carboxymethyl Cellulose with calcium chloride 0.5% for 5 minutes. The highest value of weight loss percentage was recorded for pods with untreated control. These results were achieved in the two growing seasons and were consistent with (Shebis *et al.*, 2022) who found that fruits treated concentrations of Carboxymethyl Cellulose with some edible coating the treatments gave less weight loss when compared to the untreated fruit as a control.

Also, oligosaccharides cause as elicitors signals inducing a defense response including wound healing. Yu *et al.*, 2018,

which might explain their effect on the weight loss reduction.

The effect of the interaction between both the storage period and the treatments on weight loss was significant. There was a gradual increase in weight loss in the pods in each treatment at the end of the storage period. From the above observations, the decrease in weight loss was very sharp after 20 days of storage for all treated plants.

2.1. Decay score:-

The data presented in Table (1) indicate that the treatment has an effect on the degree of decomposition of green bean pods during the storage period. The data show that the decomposition began at a lower rate and increased with the lengthening of the storage period in both seasons. The explanation for this phenomenon may be that it occurred as a result of the continuous biochemical changes that occurred in the centuries (Shebis *et al.*, 2022).

For treatments used were recorded lowest decay score in comparison to control. Moreover, green beans pods treated for Carboxymethyl Cellulose with gum arabic 5 % for 5 minutes or Carboxymethyl Cellulose with calcium chloride 0.5% for 5 minutes were the most effective treatments in reducing the decay. The highest value of decay score was recorded with untreated control.

Calcium chloride and gum arabic are used to delay and slow down the ripening of fruits and maintain their quality during the storage period. These results are in agreement with Abdullah *et al.*, 2018b.

As for the interaction between the treatments and the storage period, the data indicated that the untreated fruits began to appear decay after only 4 days of storage and many symptoms of decay were observed at the end of the storage period, while no decay was observed in the pods treated with the used materials (Carboxymethyl Cellulose with gum Arabic) till 16 days in both season.

2.2. General appearance score:-

By looking at the data in Table (1), the effect of post-harvest treatments on the general appearance (grade) of green bean pods during storage is clear. The results indicated a significant decrease in the general appearance of green bean pods with an extended storage period in both seasons. The decrease of general appearance during the storage period and reaches its maximum depression at 20 days after storage. Similar results were reported by Kasim and Kasim, 2015 and Abdullah *et al.*, 2018a on fresh-cut green bean.

In view of the effect of post-harvest treatments, the data showed clear significant differences between the treated and untreated fruits (control). Green bean pods treated with all treatments recorded clear significant differences. However, green bean pods treated with carboxymethyl cellulose with gum arabic 5% were the most effective treatments to preserve the general shape of the fruits, followed by carboxymethyl cellulose with calcium chloride 0.5%, while the untreated control recorded the least effective treatment. These results were recorded in the two seasons.

The effect of gum arabic and calcium chloride on maintaining the general appearance might be referred to its ability on decreasing fruit water loss (Abdullah *et al.*, 2018b). As for the interaction between the storage period and post-harvest treatments, the data indicated that the pods dipped in a solution of carboxymethyl cellulose with gum arabic did not have any changes in their appearance until 16 days of storage of the fruits. On the other hand, untreated

Pods had the poorest data for pod appearance at the end of the storage period (20 days). These results were consistent in both seasons.

2.3. Firmness

The data presented in Table (2) shows the effect of green bean pod firmness with post-harvest treatments during storage. Examining the data showed that there is a negative effect on a significant decrease in the firmness of green bean pods by increasing the storage period in the two seasons. The decrease in firmness during storage may be due to the activity of physiological processes within the plant, including the metabolism process, which may increase the activity of fungi. Similar results were obtained by Kassem and Kassem (2015).

For the effect of postharvest treatments, data showed that all postharvest treatments had a significant effect on pod firmness as compared with untreated control during storage. However, green bean pods treated for carboxymethyl cellulose with gum arabic 5% gave the highest value of pod firmness during storage followed by carboxymethyl cellulose with calcium chloride 0.5%. The lowest value of pod firmness was obtained from untreated control. These results were achieved in the two seasons and were in agreement with Shebis *et al.*, 2022.

The effect of the interaction between the storage period and post-harvest treatments had a significant effect on the firmness of pods in both seasons. The pods from all the treated treatments over the storage period were much better than the untreated control.

2.4. Total soluble solids percentage

The data in Table (2) shows the effect of green bean pods using post-harvest treatments on TSS as a percentage during storage. The data indicated that there was a decrease in the total dissolved solids of green bean pods in the case of prolonging the storage period until the end of the storage period in both seasons, and these results are similar Abdullah *et al.*, 2018a.

Regarding the effect of post-harvest treatments, the data showed the extent to which there was a difference in the percentage of TSS between the treated and untreated control during storage period. Green bean pods treated with carboxymethyl cellulose with gum arabic recorded the highest percentage of dissolved solids, followed by carboxymethyl cellulose with calcium chloride. Pods from all treatments during the storage period were higher in T.S.S. of those untreated control fruits. Moreover, the pods treated with carboxymethyl cellulose with gum arabic during the storage period had slightly higher TSS compared with other treatments during the storage period.

2.5. Chlorophyll content

Data in Table (2) show the effect of postharvest treatments on chlorophyll content of green bean pods during storage. Data presented that progressive and significant decrease happened in chlorophyll content. Generally, green bean pods became yellowish green in the end of the storage period. This decrement in content of chlorophyll could be attributed to the destruction by turn of chloroplasts to chromoplasts and chlorophyllase activity. These results were true in both seasons and were in agreement with Abed El-Hakim (2017) on fresh-cut Green Bean Pods.

Regarding the effect of postharvest treatments, it is clear from the results that pods treated for carboxymethyl cellulose with gum arabic significantly reduced chlorophyll degradation of green bean pods compared with other treatments, in both seasons. In general, untreated green

bean pods (control) contained the lowest value in chlorophyll contents. These results were in agreement with the findings of Shebis *et al.* (2022).

The effect of the interaction between storage period and post-harvest treatments on pods was a significant reading of chlorophyll in all planting seasons. However, pods treated with carboxymethyl cellulose and gum arabic had the highest chlorophyll content during cold storage. On the other hand, untreated control fruits recorded the lowest value of chlorophyll content during cold storage.

2.6. Total phenolic compound:

The data presented in Table (3) show the effect of total phenolic compounds content of green bean pods as a result of post-harvest treatments on storage. The data indicated a decrease in the total phenolic compound of green bean pods when the storage period was extended until the end of the storage period in the two study seasons. These results are similar to Abdul Hakim, 2017.

Concerning the effect of postharvest treatments, data indicated that there were significant differences between all treatments and clearly observed that all treatments maintained phenolic compounds during storage period with little changes in their contents, especially for treated carboxymethyl cellulose with gum arabic which show the lowest decreasing rates followed by carboxymethyl cellulose with calcium chloride compared with untreated green bean pods which had a significantly greater decrease. These results may be due to coating materials gum arabic and calcium chloride, which may form a protective barrier on the pods surface and reduce the oxygen supply for enzymatic oxidation of phenols. The same results were in agreement with Macheix *et al.*, 1990; Feng *et al.*, 2015.

By studying the effect of interaction between all post-harvest treatments and storage period on pods was significant for total phenolic compound in both seasons. However, pods treated for carboxymethyl cellulose with gum arabic was the most effective in maintaining total phenolic compound during cold storage. On the other hand, untreated control recorded lowest value of total phenolic compound during cold storage.

2.7. Microbial analysis

As shown in Table (3), results indicate that there was a linear relation between the microbial load and storage period, where it was found that the increase in storage period was accompanied by an increase in microbial load these results are similar with Abdullah *et al.*, 2018a.

Regarding the effect of postharvest treatments, it is clear from the results that the most effective antimicrobial treatment which diminishes the microbial load was treated green pods for carboxymethyl cellulose with gum arabic followed by carboxymethyl cellulose with calcium chloride compared with untreated green bean pods. Same result was obtained by Shebis *et al.*, 2022 who found that treated green bean pods with oligo carboxymethyl cellulose decrease the total bacterial counts.

Concerning the interaction among different treatments and storage period, there was a significant difference between them, where it was noticed that all treatments affected the microbial load and decrease it until the first 4 days compared with untreated pods especially green pods treated for carboxymethyl cellulose with gum arabic. Then after this storage period a remarkable increase in microbial load was observed.

Table (1): Effect of carboxymethyl cellulose with edible coating treatments on weight loss %, decay and general appearance of green beans during cold storage.

Treatments	Days after storage	First season			Second season		
		Weight loss %	Decay	General appearance	Weight loss %	Decay	General appearance
Control	0		1.00 g	9.00 a		1.00 f	9.00 a
	4	1.45 q	1.33 fg	8.33 ab	1.07 t	1.33 ef	8.33 ab
	8	1.95 n	2.00 de	7.00 cd	1.54 o	1.67 de	7.00 cd
	12	3.17 g	2.33 cd	6.33 d	2.94 h	2.00 cd	6.33 de
	16	3.95 c	3.00 ab	5.00 e	3.54 c	2.67 ab	5.00 f
CMC+ Gum Arabic	0		1.00 g	9.00 a		1.00 f	9.00 a
	4	0.93 w	1.00 g	9.00 a	0.75 w	1.00 f	9.00 a
	8	1.25 t	1.00 g	9.00 a	1.15 s	1.00 f	9.00 a
	12	2.16 m	1.00 g	9.00 a	1.94 n	1.00 f	9.00 a
	16	2.84 i	1.00 g	8.33 ab	2.65 j	1.00 f	8.33 ab
CMC+ CaCl ₂	0		1.00 g	9.00 a		1.00 f	9.00 a
	4	1.11 v	1.00 g	9.00 a	0.82 v	1.00 f	9.00 a
	8	1.38 r	1.00 g	9.00 a	1.28 r	1.00 f	9.00 a
	12	2.35 l	1.00 g	9.00 a	2.23 m	1.00 f	9.00 a
	16	3.13 h	1.33 fg	7.00 cd	2.83 i	1.33 ef	7.00 cd
CMC	0		1.00 g	9.00 a		1.00 f	9.00 a
	4	1.33 s	1.00 g	9.00 a	0.83 v	1.00 f	9.00 a
	8	1.67 o	1.00 g	9.00 a	1.43 p	1.00 f	9.00 a
	12	2.76 j	1.33 fg	8.33 ab	2.55 k	1.00 f	9.00 a
	16	3.65 d	2.00 de	6.33 d	3.13 f	1.67 de	6.33 de
CMC+ Ascorbic acid	0		1.00 g	9.00 a		1.00 f	9.00 a
	4	1.19 u	1.00 g	9.00 a	0.88 u	1.00 f	9.00 a
	8	1.56 p	1.00 g	9.00 a	1.33 q	1.00 f	9.00 a
	12	2.55 k	1.00 g	9.00 a	2.41 l	1.67 de	7.00 cd
	16	3.44 e	1.67 ef	6.33 d	2.94 h	1.67 de	6.33 de
Control		3.03 A	2.17 A	6.67 C	2.69 A	1.94 A	6.44 D
CMC+ Gum Arabic		2.10 E	1.06 D	8.67 A	1.91 E	1.06 D	8.67 A
CMC+ CaCl ₂		2.32 D	1.22 CD	8.00 B	2.08 D	1.17 CD	8.11 B
CMC		2.70 B	1.50 B	7.67 B	2.31 B	1.33 BC	7.67 C
CMC+ Ascorbic acid		2.53 C	1.28 C	7.89 B	2.20 C	1.39 B	7.56 C
	0		1.00 D	9.00 A		1.00 D	9.00 A
	4	1.20 E	1.07 D	8.87 A	0.87 E	1.07 D	8.87 A
	8	1.56 D	1.20 CD	8.60 AB	1.35 D	1.13 CD	8.60 A
	12	2.60 C	1.33 C	8.33 B	2.42 C	1.33 C	8.07 B
	16	3.40 B	1.80 B	6.60 C	3.02 B	1.67 B	6.60 C
	20	3.93 A	2.27 A	5.27 D	3.53 A	2.07 A	5.00 D

Values followed by the same letter (s) are not significantly different at 5 %

Table (2): Effect of carboxymethyl cellulose with edible coating treatments on firmness (kg/cm²), total soluble solids % and total chlorophyll of green beans during cold storage.

Treatments	Days after storage	First season			Second season		
		Firmness (kg/cm ²)	Total soluble solids %	Total chlorophyll	Firmness (kg/cm ²)	Total soluble solids %	Total chlorophyll
Control	0	2.66 a	6.40 a	26.31 a	2.57 a	6.00 a	25.66 a
	4	2.24 g	6.20 bcd	25.31 h	1.78 fgh	6.00 a	24.72 f
	8	1.73 m	6.27 abc	25.05 k	1.62 hi	5.87 bc	24.03 l
	12	1.44 p	6.07 def	24.63 o	1.22 lm	5.73 de	23.74 n
	16	1.03 t	5.87 ghi	23.82 u	0.88 no	5.47 h	23.13 r
	20	0.73 u	5.73 i	23.24 v	0.92 n	5.47 h	22.54 t
CMC+ Gum Arabic	0	2.66 a	6.40 a	26.31 a	2.57 a	6.00 a	25.66 a
	4	2.53 b	6.27 abc	25.91 b	2.36 b	6.00 a	25.30 b
	8	2.36 d	6.33 ab	25.60 e	2.11 cd	5.93 ab	25.01 d
	12	2.13 h	6.20 bcd	25.24 j	1.86 efg	5.80 cd	24.61 g
	16	1.80 l	6.00 efg	24.81 n	1.61 hi	5.67 ef	24.22 j
	20	1.42 pq	5.80 hi	24.36 r	1.27 kl	5.60 fg	23.83 m
CMC+ CaCl ₂	0	2.66 a	6.40 a	26.31 a	2.57 a	6.00 a	25.66 a
	4	2.41 c	6.27 abc	25.78 c	2.30 b	6.00 a	25.11 c
	8	2.23 g	6.27 abc	25.36 g	2.02 de	5.87 bc	24.72 f
	12	1.93 j	6.07 def	25.02 l	1.70 gh	5.80 cd	24.23 j
	16	1.64 n	6.00 efg	24.54 p	1.41 jk	5.60 fg	23.82 m
	20	1.22 s	5.73 i	24.02 t	1.05 mn	5.47 h	23.52 p
CMC	0	2.66 a	6.40 a	26.31 a	2.57 a	6.00 a	25.66 a
	4	2.30 f	6.20 bcd	25.60 e	2.25 bc	6.00 a	24.93 e
	8	1.91 jk	6.13 cde	25.28 i	1.81 fg	5.87 bc	24.44 h
	12	1.61 o	6.07 def	24.91 m	1.51 ij	5.67 ef	24.14 k
	16	1.28 r	5.87 ghi	24.17 s	1.13 lm	5.53 gh	23.56 o
	20	1.90 k	5.73 i	23.81 u	0.73 o	5.47 h	23.05 s
CMC+ Ascorbic acid	0	2.66 a	6.40 a	26.31 a	2.57 a	6.00 a	25.66 a
	4	2.33 e	6.27 abc	25.71 d	2.27 bc	6.00 a	25.11 c
	8	2.04 i	6.20 bcd	25.41 f	1.92 ef	5.87 bc	24.74 f
	12	1.73 m	6.07 def	25.02 l	1.62 hi	5.67 ef	24.30 i
	16	1.40 q	5.93 fgh	24.50 q	1.29 kl	5.60 fg	23.84 m
	20	1.02 t	5.73 i	24.02 t	0.88 no	5.53 gh	23.27 q
Control		1.64 E	6.09 B	24.73 D	1.50 E	5.76 B	23.97 E
CMC+ Gum Arabic		2.15 A	6.17 A	25.37 A	1.96 A	5.83 A	24.77 A
CMC+ CaCl ₂		2.02 B	6.12 AB	25.17 B	1.84 B	5.79 AB	24.51 B
CMC		1.94 C	6.07 B	25.01 C	1.67 D	5.76 B	24.30 D
CMC+ Ascorbic acid		1.86 D	6.10 B	25.16 B	1.76 C	5.78 B	24.49 C
	0	2.66 A	6.40 A	26.31 A	2.57 A	6.00 A	25.66 A
	4	2.36 B	6.24 B	25.66 B	2.19 B	6.00 A	25.04 B
	8	2.06 C	6.24 B	25.34 C	1.90 C	5.88 B	24.59 C
	12	1.77 D	6.09 C	24.97 D	1.58 D	5.73 C	24.20 D
	16	1.43 E	5.93 D	24.37 E	1.26 E	5.57 D	23.72 E
	20	1.26 F	5.75 E	23.89 F	0.97 F	5.51 E	23.24 F

Values followed by the same letter (s) are not significantly different at 5 %

Table (3): Effect of carboxymethyl cellulose with edible coating treatments on total phenolic (mg/100 g) and microbial analysis of green beans during cold storage.

Treatments	Days after storage	First season				Second season			
		Total phenolic mg/100 g		Microbial analysis		Total phenolic mg/100 g		Microbial analysis	
Control	0	153.13	a	0.33	w	151.19	a	0.31	l
	4	148.34	e	1.34	s	144.52	f	1.26	ijk
	8	140.65	j	2.54	k	137.41	j	2.55	ef
	12	134.23	m	3.32	f	127.72	r	3.84	bc
	16	126.03	r	4.14	c	118.54	w	3.93	b
	20	119.12	s	4.95	a	101.35	x	6.76	a
CMC+ Gum Arabic	0	153.13	a	0.33	w	151.19	a	0.31	l
	4	151.11	b	0.93	v	148.84	b	0.85	k
	8	147.51	f	1.54	r	145.02	e	1.45	hij
	12	142.83	h	1.95	o	138.17	i	1.95	gh
	16	137.92	k	2.75	i	132.72	m	2.54	ef
	20	132.52	n	3.23	g	129.12	p	3.03	de
CMC+ CaCl ₂	0	153.13	a	0.33	w	151.19	a	0.31	l
	4	150.91	bc	1.04	u	147.02	c	0.95	jk
	8	147.50	f	1.75	q	143.18	g	1.75	ghi
	12	141.52	i	2.23	m	135.15	k	2.24	fg
	16	136.52	l	2.96	h	130.43	o	2.82	e
	20	130.71	p	3.85	d	126.73	s	3.35	cd
CMC	0	153.13	a	0.33	w	151.19	a	0.31	l
	4	149.85	d	1.08	t	146.42	d	1.11	jk
	8	143.13	h	2.13	n	141.72	h	2.17	fg
	12	138.03	k	2.63	j	131.84	n	2.85	de
	16	131.35	o	3.53	e	125.62	t	3.36	cd
	20	127.63	q	4.43	b	121.23	v	4.03	b
CMC+ Ascorbic acid	0	153.13	a	0.33	w	151.19	a	0.31	l
	4	150.63	c	1.09	t	147.03	c	1.01	jk
	8	145.32	g	1.92	p	143.33	g	1.93	gh
	12	140.42	j	2.45	l	133.24	l	2.55	ef
	16	134.12	m	3.34	f	128.14	q	3.05	de
	20	130.55	p	4.17	c	124.52	u	3.75	bc
Control		136.91	E	2.77	A	130.12	E	3.11	A
CMC+ Gum Arabic		144.17	A	1.79	E	140.84	A	1.69	D
CMC+ CaCl ₂		143.38	B	2.03	D	138.95	B	1.90	CD
CMC		140.52	D	2.36	B	136.34	D	2.31	B
CMC+ Ascorbic acid		142.36	C	2.21	C	137.91	C	2.10	BC
	0	153.13	A	0.33	F	151.19	A	0.31	F
	4	150.17	B	1.10	E	146.77	B	1.04	E
	8	144.82	C	1.98	D	142.13	C	1.97	D
	12	139.41	D	2.52	C	133.22	D	2.69	C
	16	133.19	E	3.34	B	127.09	E	3.14	B
	20	128.10	F	4.12	A	120.59	F	4.18	A

Values followed by the same letter (s) are not significantly different at 5 %

4. References:

[1] O. A. C., *Official methods of analysis of AOAC International*. 17th edition. Gaithersburg, MD,

USA, Association of Analytical Communities (2000).

[2] Abed EL-Hakim, H. I., Improvement of Shelf Life and Physicochemical Quality of Fresh-Cut

- Green Bean Pods (*Phaseolus vulgaris* L. cv. Polista) Using Edible Coatings. *Journal of Food Sciences*; Suez Canal University, 4(1) (2017) 1-10.
- [3] Adetunji, C. O., Arowora, K.A., Fawole O. B., Adetunji, J. B. and Olagbaju, A. R., Effect of edible coating of carboxymethyl cellulose and corn starch on cucumber stored at ambient temperature. *Asian Journal of Agriculture and Biology*, 1(3) (2013) 133-140.
- [4] Ali, A., Maqbool, M., Ramachandran, S. and Alderson, P. G., Gum arabic as a novel edible coating for enhancing shelf-life and improving postharvest quality of tomato (*Solanum lycopersicum* L.) fruit. *Postharvest Biology and Technology*, 58 (2010) 42–47.
- [5] Al-Sanabani, A. S., K. M. Youssef, A. A. Shatta and S. K. El-Samahy, Influence of freezing steps in color attributes phytochemical contents and antioxidant capacity of green beans. *J. Food Sci.*, Suez Canal Univ., 3(1) (2016) 19- 25.
- [6] Anderson, D. M. W. and Eastwood, M. A., The safety of gum arabic as a food additive and its energy value as an ingredient: a brief review. *Journal of Human Nutrition and Dietetics*, 2(1989)137–144.
- [7] Anese, M. M. and Nicoli, M. C., Quality of minimally processed apple slice using different modified atmosphere conditions. *J. Food Quality*, 20 (1997) 359 – 370.
- [8] Bilgin, J., The Effects of Salicylic, Folic and Ascorbic Acid Treatment on Shelf Life Quality of Broccoli Florets. *Journal of Agricultural Production*, 2(1) (2021) 7-15. doi: 10.29329/agripro.2021.344.2 *Bioscience Research*, 15(4):3625-3634.
- [9] Biswal, D. R. and Singh, R. P., Characterisation of carboxymethyl cellulose and polyacrylamide graft copolymer. *Carbohydrate polymers*. 57 (2004) 379-387.
- [10] Duncan, D. B., 1955. Multiple range and multiple F test. *J. Biometrics*, 11: 1-42.
- [11] El-Anany, A. M., Hassan, G. F. A. and Rehab Ali, F.M., Effects of edible coatings on the shelf-life and quality of Anna apple (*Malus domestica Borkh*) during cold storage. *Journal of Food Technology* 7 (2009) 5–11.
- [12] El-Mogy, M. M.; Parmar, A., Ali, M. R., Abdel-Aziz, M. E. and Abdeldaym, E. A., Improving postharvest storage of fresh artichoke bottoms by an edible coating of Cordia myxa gum. *Postharvest Biol. Technol.*, 163 (2020)111143.
- [13] Fenet Gameda, Effect of Post-Harvest Calcium Chloride Treatment on Quality and Shelf Life of Apple (*Malus domestica*). *International Journal of Food Engineering and Technology*. 5(2) (2021) 62-67. doi: 10.11648/j.ijfet.20210502.16
- [14] Feng, D., L. Shujun, L. Zhongming, Z. Kexin, W. Xiaolin and J. Chunde, Improvement of quality and shelf life of strawberry with nanocellulose/chitosan composite coatings. *Bangladesh J. Bot.*, 44(5) (2015) 709-717.
- [15] Huang Q., Wan C., Zhang Y., Chen C. and Chen J., Gum Arabic Edible Coating Reduces Postharvest Decay and Alleviates Nutritional Quality Deterioration of Ponkan Fruit During Cold Storage. *Front. Nutr.* 8 (2021) 717596. doi: 10.3389/fnut.2021.71759.
- [16] Jimenez, M., E. Trejo and M. Cantwell, Postharvest quality changes in green beans. *Research Report*, UC Davis, cooperative extension service No. pp9 (1998).
- [17] Kasim, R. and M. U. Kasim, Biochemical changes and color properties of fresh-cut green bean (*Phaseolus Vulgaris* L.cv. gina) treated with calcium chloride during storage. *Food Sci. Technol. Campinas*, 35(2) (2015) 266-272.
- [18] Kasinee S., Rujira D., and Lampan K., Effect of Carboxymethyl Cellulose as Edible Coating on Postharvest Quality of Rambutan Fruit under Ambient Temperature. *International Journal of Agricultural Technology*, 13(7.1) (2017) 1449-1457.
- [19] Macheix, J. J., A. Fleuriet and J. Bilot (EDS.), *Fruit phenolics*. CRC Press, Inc., Florida, (1990).
- [20] Maftoonazad, N. and Ramaswamy, H. S., 2005. Postharvest shelf -life extension of avocados using methylcellulose-based coating. *LWT-Food Science and Technology*, 38:617–624.
- [21] Mohamed A. A. Abdullah, Huda A. Ibrahim, Nagwa M. K. Hassan and Heba S. El-Batran, Use of Gum arabic and Calcium Chloride as Edible Coating to Delay Postharvest Ripening and to Maintain Melon Quality during Storage. *Bioscience Research*. 15(4) (2018b) 3625-3634.
- [22] Mohamed A. A. Abdullah, Ramadan A. H. and Amr M. M., Effect of Gamma Irradiated Chitosan and Green Tea on Pod Quality of Fresh-Cut Green Bean under Cold Storage Conditions. *Journal of Horticultural Science & Ornamental Plants*, 10(3) (2018a) 140-149.
- [23] Motlagh, S., Ravines, P. and Karamallah, K. A., Ma, Q., The analysis of Acacia gums using electrophoresis. *Food Hydrocolloid*, 20 (2006) 848–854.
- [24] Naveena, B. and Immanuel, G., Effect of Calcium Chloride, Sodium Chloride and Lime Juice on PhysicoChemical Properties of Cucumber. *International Journal of Agricultural Science and Research (IJASR)*, 7 (4) (2017) 765-770.
- [25] Rachtanapun, P. and Rattanapanone, N., Synthesis and characterization of carboxymethyl cellulose powder and films from *Mimosa pigra*. *Journal of Applied Polymer Science*, 122 (2011) 3218–3226.
- [26] Rachtanapun, P., Kumthai, S., Yagi, N. and Uthaiyod, R., Production of carboxymethyl cellulose (CMC) films from papaya peel and their mechanical properties. *Proceedings of the Kasetsart University Annual Conference*:

- Agricultural Extension and Home Economics: Agro-Industry, (2007) 790 -799.
- [27] Rachtanapun, P., Luangkamin, S., Tanprasert, K. and Suriyatam, R., Synthesis and characterization of carboxymethyl cellulose from durian rind. Proceedings of 49th Kasetsart University Annual Conference: Agro-Industry, (2011).
- [28] Rachtanapun, P., Thanakkasane, S. and Soonthornampai, S., Application of Carboxymethyl cellulose from Papaya Peel for Mango (*Mangifera indica* L.) 'Namdokmai' Coating. Journal of Agricultural Science, 39(3) (2008) 74-82.
- [29] Rójas-Graü, M. A., R. Soliva-Fortuny and Q. MortisBelloso, Using of edible coating for fresh-cut fruits and vegetables. In: Advances in fresh-cut fruits and vegetables processing. (Mortis-Belloso, Q. and Soliva-Fortuny, R. eds.), Chll., (2011) 285-304. CRC press Taylor and Francis Group. Boca Ratan, London, New York.
- [30] Saleh, S. A., Liu, G., Liu, M., Ji, Y., He, H., and Gruda, N., Effect of irrigation on growth, yield, and chemical composition of two green bean cultivars. Horticulturae (2018) 4, 3. doi: [10.3390/horticulturae4010003](https://doi.org/10.3390/horticulturae4010003)
- [31] Shalata, A. and P. M. Neumann, Exogenous ascorbic acid (vitamin C) increases resistance to salt stress and reduced lipid peroxidation. Experimental botany, 52(346) (2001) 2207-2211.
- [32] Shebis, Y., Fallik, E., Rodov, V., Sagiri, S.S. and Poverenov, E., Oligomers of Carboxymethyl Cellulose for Postharvest Treatment of Fresh Produce: The Effect on Fresh-Cut Strawberry in Combination with Natural Active Agents. Foods (2022) 1108- 1117. <https://doi.org/10.3390/foods11081117>
- [33] Shehata, S.A., Elmogy, M. and Mohamed, H.F.Y., Postharvest quality and nutrient contents of long sweet pepper enhanced by supplementary potassium foliar application. Int. J. Veg. Sci., 25 (2018) 196–209.
- [34] Snedecor, C. W., and Cochran, W. G., Statistical Methods. 7th Ed. The Iowa State Univ. Press. Ames, Iowa, USA, (1982) 325-330.
- [35] Toğrul, H. and Arslan, N., Extending shelf-life of peach and pear by using CMC from sugar beet pulp cellulose as a hydrophilic polymer in emulsions. Food Hydrocolloids, 18 (2004) 215-226.
- [36] Watada, A. E. and L. L. Morris, Effect of chilling and non-chilling temperatures on snap beans fruits. Proc. Amer. Soc. Hort. Sci., 89 (1996) 368-374.
- [37] Yu, Y., Gim, S., Kim, D.; Arnon, Z., Gazit, E., Seeberger, P.H. and Delbianco, M., Oligosaccharides Self-Assemble and Show Intrinsic Optical Properties. J. Am. Chem. Soc., 141 (2019) 4833–4838.
- [38] Zhu, X., Wang, Q., Cao, J., Jiang, W., Effects of chitosan coating on postharvest quality of mango (*Mangifera indica* L: cv. Tainong) fruits. J. Food Process. Preserv., 32 (2008) 770–784.