



Inflammatory Cytokines, Oxidative Stress Biomarkers and Clinical Manifestations of Organophosphorus Pesticides-Exposed Researchers

Nermeen Said Abdel Aziz^a, Mohamed Yehia Elawady^b, Sanaa Aboulmakarem Rizk^a, Sally Adel Hakim^b, Eman Mohamed Shahy^a, Ebtessam Abdel-Shafy^a



CrossMark

^a Department of Environmental and Occupational Medicine, Division of Environmental Research, National Research Centre
^b Public Health and Occupational Medicine Department, Faculty of Medicine, Ain Shams University

Abstract

Organophosphorus pesticides (OPs) exposure may be associated with an increased risk of immunotoxicity in the occupational workplace. The study was designed to investigate and compare the correlation between clinical symptoms, oxidative stress, and the immunological effects of OPs exposure among occupationally exposed researchers. Clinical manifestations associated with pesticide exposure were collected from 49 researchers who were occupationally exposed to OPs (exposed group), and 51 non-exposed researchers (control group). The levels of cholinesterase enzymes (Acetylcholinesterase (AChE) and Butyrylcholinesterase (BuChE)) have been estimated as an exposure biomarker for pesticide exposure. The oxidative stress status was evaluated by malondialdehyde (MDA) assessment, while total antioxidant capacity (TAC) could be an indicator of the antioxidant mechanism. We used Interleukin -6 (IL-6) and Interferon- gamma (IFN- γ) as immune biomarkers as well as Tumor Necrosis factor - Alpha (TNF- α) as a biomarker of cellular immune function. Our results revealed a significant decline in AChE, BuChE, and TAC levels, while there was a significant elevation in MDA in occupationally exposed researchers versus control group. In addition, the levels of IL-6, IFN- γ , and TNF- α have been increased in exposed researchers. From clinical examination, the chief morbidity was pertained to neurological disorders and bronchial asthma as a result of occupational exposure to OPs pesticides. Muscular weakness was significantly syndicated with the decline in AChE and BuChE levels. Bronchial asthma was closely associated with MDA, IL 6, and TNF- α . Furthermore, in numbness, MDA, IL-6, IFN- γ , and TNF- α , were significantly higher. This study also indicated that MDA, IL-6, and TNF- α levels were significantly boosted up in exposed researchers who were not wearing a mask. This could be a risk factor for some adverse health hazards.

Keywords: Organophosphorus Pesticides, AChE, BuChE, Cytokines, oxidative stress, MDA, TAC, Personal Protective Equipment (PPE)

Introduction

Organophosphorus are compounds that most commonly used in agriculture as pesticides¹. Despite its popularity and widespread prevalence, the pesticides raise serious concerns about health risks arising from prolonged exposure². Occupational exposure to OPs may cause immunotoxicity, cardiovascular, neurological, reproductive and carcinogenic disorders³. There are two types of cholinesterase enzymes; acetylcholinesterase (AChE) that present in neural tissues and the erythrocyte membrane⁴ and plasma cholinesterase (BuChE) that is synthesized primarily

in the liver and released at a high rate into plasma⁵. Prolonged exposure to OPs ceases cholinesterase activities. However, cholinesterase inhibition alone cannot explain the wide range of clinical disorders⁶. In addition, Oxidative stress and disturbances of cytokine balance are other factors that contribute to pesticides health hazards⁶. Oxidative stress-induced by pesticides enhances reactive oxygen species (ROS) that alter the oxidative balance within cells⁷. Antioxidants considered as scavenging enzymes act as defense mechanisms in the detoxification of free radicals⁷. Severity of oxidative damage can be estimated by measuring malondialdehyde (MDA, the end product of lipid peroxidation), while the

*Corresponding author e-mail: zahran_nermeen@yahoo.com; (Nermeen Zahran).

Receive Date: 09 December 2020, Revise Date: 10 January 2021, Accept Date: 10 January 2021

DOI: 10.21608/EJCHEM.2021.52948.3094

©2021 National Information and Documentation Center (NIDOC)

total antioxidant capacity (TAC) is one of the antioxidant defense mechanisms⁸. As the balance between pro-inflammatory and anti-inflammatory mediators regulate the inflammatory response⁹ and OPs induce low-grade inflammation, so, chronic exposure to OPs cause disturbances between the release of pro-inflammatory mediators¹⁰. Cytokines such as Tumor Necrosis factor – Alpha (TNF- α), interleukin (IL-1, IL-6 and IL-10), or type 1 interferon (IFN) secretion increased by activated macrophages that leads to up-regulation of inflammatory reactions¹⁰, and anti-inflammatory cytokines, that inhibit pro-inflammatory cytokine synthesis and adhesion molecule expression while increasing the levels of specific cytokine inhibitors⁹. Therefore, it was useful to investigate the disturbance of the cytokine balance from exposure to pesticides. Occupational exposures to pesticides are mainly due to inhalation and skin absorption¹¹. Respiratory inhalation occurs when OPs are sprayed in "an inhalable form". Dermal absorption occurs through skin contact with pesticides or through clothing and tools contaminated with pesticide residues¹¹.

The goal of this study was therefore to assess the association between the inflammatory cytokines and the oxidative stress biomarkers with clinical manifestations and their correlation with the use of PPE in researchers who are occupationally exposed to pesticides.

2. Subjects and Methods and sampling

2.1. Study Design and Population

A cross-sectional study was conducted in 49 researchers occupationally exposed to pesticides and 51 non-occupationally exposed controls. Both groups were recruited from different Egyptian governmental research centers, and matched their socioeconomic status, age, and smoking habits. The approval of the Ethics Committee was received from the Ethical Committee at the National Research Centre, Egypt. Written consent was obtained from all the selected subjects.

2.2. Questionnaire & Clinical examination

The questionnaire included details of the socio-demographic aspects, smoking habits, and history of health problems for any system of the body. Also, clinical examination of all systems was performed with special concern of respiratory and neurological systems.

2.3 Sample collection

About 5 ml of the venous blood was collected from each subject and divided into two separate tubes. Two ml of blood was left in a dry tube for 30 minutes at room temperature for coagulation and then centrifuged for 10 minutes at 3000 rpm for serum separation for AChE, TAC, MDA, IL6, IFN- γ and TNF- α determination. A further three ml of blood was collected in an EDTA tube, centrifuged for 3000 rpm for 10 minutes to separate plasma for determination of BuChE. All samples were stored at -20° C until determination by colorimetric method.

2.4 Determination of cholinesterase activities

2.4.1 Acetylcholinesterase

The immunoassay kit was used to measure the activity of human acetylcholinesterase in serum in vitro by (Sandwich-ELISA, Sunredbio, Shanghai).

2.4.2 Butyrylcholinesterase

The activity of butyrylcholinesterase was measured in plasma using a colorimetric method according to Knedel and Bottger¹².

2.5 Determination of oxidative stress markers

2.5.1 Malondialdehyde

Serum MDA activity was colorimetrically determined in the serum¹³.

2.5.2 Total antioxidant capacity

This assay based on the ability of the enzyme to inhibit the phenazine methosulphate -mediated reduction of nitroblue tetrazolium dye.

2.6 Immunological Profile

2.5.3 Serum Cytokines

Cytokines (IL-6, TNF- α and IFN- γ) were quantitatively analyzed using ELISA kit purchased from SinoGene Biotech Co.,Ltd, China.

3- Statistical analysis

Statistical analysis of the collected data and the laboratory results was performed with SPSS version 24. The comparisons of quantitative variable between two groups were performed using independent t-test and expressed as mean and standard deviation (Mean \pm SD). Qualitative results were expressed as number (no.) and percent (%). Qualitative results were analyzed using

Pearson's Chi-square (χ^2) and likelihood ratio will be used in cases where more than 25% of the cells had expected count < 5 . Pearson correlation coefficient was used to study the relationship between the different quantitative variables.

4-Results

The exposed researchers were occupationally exposed to OPs for about 6.3 ± 1.9 years. There was no significant difference in age (P -value=0.38) between workers and control groups (39.4 ± 1.42 and 37.02 ± 1.79 respectively). The exposed researchers consisted of 38(77.6%) males and 11(22.4%) females while the control group contained 35 (68.6%) males and 16 (31.4%) females. There was no significant difference in male and female percentage between two groups (P -value =0.371). About 7 (14.3%) of the exposed workers were smokers while 42 (85.7%) were not smokers. In contrast, 11 (21.6%) of the controls were smokers while 40 (78.4%) were not. There was no significant difference in the smoking habits (P -value =0.289) between the two groups.

By means of a questionnaire and clinical examination, several clinical risks were detected in exposed populations. Fig. (1) represents different clinical manifestations; neurological disorders (muscle weakness (53.06% and numbness 59.18%), respiratory dysfunction (bronchial asthma 51.02%), eye irritation (20.41%), and dermatitis (30.61%).

Concerning estimated biochemical parameters;

these results indicated a significant decline in biomarkers of exposure (AChE and BuChE) in exposed researchers relative to the control group. There was a significant elevation in oxidative stress biomarker (MDA), inflammatory cytokines (IL-6, IFN- γ , and TNF- α) while there was a significant decline in the TAC level.

Fig.2 represents a significant negative correlation between duration of exposure and BuChE ($r = 0.104$, p value < 0.05) while there was a significant positive correlation between duration of exposure with MDA ($r = 0.072$, p value < 0.05), IL-6 ($r = 0.678$, p value < 0.01) and TNF- α ($r = 0.544$, p value < 0.01).

Table (2) showed that in exposed researchers, AChE was positively correlated with BuChE and TAC but negatively correlated with IL-6 while BuChE was negatively correlated with IL-6 and IFN- γ . Additionally, TAC was negatively correlated IL-6 and IFN- γ . There was appositve correlation between MDA with IL-6 and TNF- α . Also, IL-6 was positively correlated with IFN- γ and TNF- α .





Figure1. Prevalence of clinical Symptoms among the exposed group.

Table 1: Comparison of different estimated biochemical parameters between researchers and their controls

Variable	Control (51)	Exposed (49)	P-value
AChE (ng /mL)	352.45±23.35	184.59±13.15	<0.001
BuChE (U/ L)	3018.06±78.33	2011.59±270.25	<0.005
TAC (mM/L)	1.23±0.05	0.70±0.02	<0.001
MDA (nmol/ml)	6.71±0.39	9.29±0.65	<0.005
IL- 6 (ng/L)	1.05±0.14	3.09±0.25	<0.001
IFN- γ (pg/ml)	663.48±84.25	1380.15±79.14	<0.001
TNF α (ng/l)	98.00±9.64	127.63±11.35	<0.005

**Figure2: The Correlation between BuChE (ng/L), MDA (nmol/ml), IL-6 (ng/L) and TNF- α (ng/l) with duration of exposure (year) in the exposed group**

Tab.2 correlation analysis between estimated biomarkers among pesticides-exposed researchers

	AChE (ng /mL)	BuChE (U/ L)	TAC (mM/L)	MDA (nmol/ml)	IL-6 (ng/L)	IFN- γ (pg/ml)
BuChE (U/ L)	0.12*	-----				
TAC (mM/L)	0.35**	0.14	-----			
MDA (nmol/ml)	-0.16	-0.13	-0.31	-----		
IL-6 (ng/L)	-0.37*	-0.32*	-0.46	0.49*	-----	
IFN-γ (pg/ml)	-0.36	-0.08*	-0.61	0.68	0.23*	-----
TNF-α (ng/l)	-0.19	-0.11	-0.28*	0.18*	0.83**	0.04

*Correlation is significant at the 0.05 level (2-tailed).

**Correlation is significant at the 0.01 level (2-tailed).

Table (3): Comparison of pesticides exposure biomarkers with clinical symptoms in the exposed group

Clinical symptoms		TAC (mM /L)	MDA (nmol/ml)	IL-6 (ng/L)	IFN- γ (pg/ml)	TNF- α (ng/l)
Muscular-weakness	with	0.51±0.02*	9.8±1.1	1532.0±90.0*	3.52±0.6	134.5±21.4
	without	0.98±0.04	8.8±0.8	1208.5±127.3	3.09±0.39	129.0±18.6
Numbness	with	0.71±0.02	11.1±1.1*	1585.6±63.0**	3.91±1.8*	134.5±16.7*
	without	0.79±0.1	8.0±0.8	1082.2±149.6	2.39±0.67	93.8±24.4
Bronchial asthma	with	0.69±0.04	17.2±5.4*	1395.9±81.4	3.60±0.63	226.12±14.1*
	without	0.73±0.01	9.0±0.6	1009.8±249.8	2.19±0.32	116.75±28.9
Eye-irritation	with	0.72±0.01	12.2±0.3*	1395.9±81.4	3.22±0.71*	131.9±16.6
	without	0.74±0.02	9.3±1.5	1391.5±87.0	1.89±0.19	115.3±20.0
Skin-dermatitis	with	0.70±0.03	8.7±0.6	1321.4±159.4*	3.31±0.41	134.5±22.0
	without	0.73±0.03	7.4±0.7	1009.8±249.8	3.09±0.41	127.9±19.9

Table (3) illustrated the association between estimated cytokines and oxidative stress biomarkers with clinical symptoms in exposed researches. Muscular weakness was combined with elevation of IL-6 levels and decline in TAC level. The levels of MDA, IL-6, IFN- γ , and TNF- α was significantly raised in researchers who represents the symptoms of numbness. In exposed subjects with bronchial asthma, there was a significant boost in MDA, IFN- γ , and TNF- α . In addition, eye-irritation was significantly related to higher levels of MDA, and IFN- γ . Skin-dermatitis was significantly linked to a

heightened IL-6 level.

Regarding, the use of PPE among exposed researchers, 61.20% were wearing gloves, 71.4% were wearing mask and 63.3% cared for wearing the apron. Our results revealed that the exposed group who was not interested in wearing gloves showed a significant decrease in BuChE and increase in TNF- α level. Respecting those who did not wear a worn mask, there was significant decrease in BuChE and significant increase in MDA, IL-6, and TNF- α levels. Exposed group who were not wearing apron demonstrated significant elevation in MDA levels.

Table (4): Comparison of biochemical estimated parameters with usage of different PPE.

parameters	PPE-gloves		PPE-mask		PPE- apron	
	Yes (71.4%)	No (28.6%)	Yes (61.2%)	No (28.8%)	Yes (63.3%)	No (36.7%)
AChE (ng /mL)	204.3±26.6	188.3±16.5	204.5±22.1	185.0±18.2	201.6±23.3	188.0±17.6
BuChE (U/ L)	3049.9±453.2	1579.3±297.01*	2682.8±386.4	1542.1±336.3*	2030.1±342.2	1981.0±422.4
TAC (mM/L)	0.72±0.02	0.70±0.02	0.71±0.02	0.68±0.03	0.73±0.04	0.71±0.02
MDA (nmol/ml)	8.11±1.34	9.52±0.69	6.21±1.21	9.01±0.84*	7.21±0.62	10.21±0.91*
IL-6 (ng/L)	2.62±0.34	3.21±0.33	2.22±0.2	3.61±0.43**	3.01±0.34	3.09±0.34
INF- γ (pg/ml)	1364.9±133.4	1313.7±106.6	1204.1±121.3	1416.1±114.2	1371.5±152.3	1303.4±100.9
TNF-α (ng/l)	77.0±2.5	146.1±14.2**	78.9±2.1	158.7±16.1**	121.8±15.0	128.2±15.2

* P-value <0.05

** P-value <0.001

5-Discussion

Although the use of pesticides has great benefits, it has several health risks associated with humans who are occupationally or environmentally exposed to these compounds¹⁴. The Occupational pesticides exposure has been previously proved to induce immune dysfunction increasing risk of several immune-associated diseases^{14,15}. The mechanism by which pesticides alter immune system homeostasis is still under discussion especially at workplace. As shown in Figure (1) there was an increase in wheezy chest and asthma in some cases of OPs exposed researchers (51.02%), and this result was in agreement with Fareed et al.¹⁶, who reported that occupational exposure to OPs is associated with an increased risk of respiratory symptoms, asthma and chronic bronchitis¹⁶. Sapbamrer et al.¹⁵, also found difficulty in breathing with chronic pesticides exposure among rice farmers occupationally exposed to OPs¹⁵. Many factors may be involved in the development of asthma related to OPs exposure such as inhibition of cholinesterase enzymes¹⁷. Drake et al.¹⁸ reported that pesticides exposure interact with AChE increasing the amount and residence time of acetylcholine at muscarinic and nicotinic receptors in target tissues, resulting in bronchoconstriction¹⁸.

Our results prevailed also the persistence of some neurological disorders in pesticides exposed

researcher; muscle weakness (53.06%) and numbness (59.19%). In consistent with our results, Sapbamrer et al.¹⁵ detected a neurological dysfunction among occupationally pesticides exposed farmers demonstrated by significant muscle weakness and numbness¹⁵. Additionally, Hongsibsong et al.¹⁹, reported muscle weakness and numbness in OPs exposed cases¹⁹. Conversely, some studies have reported no strong associations between occupational exposure and neurobehavioral performance or neurodegeneration²⁰. According to our study, Eye irritation also showed a significant increase in exposed compared to the control group (20.41%). Also, dermatitis was detected in (30.61%) of exposed researchers that was mostly in the forearm and hands with relapses and remissions. Our results agreed with Farahat et al.²¹, who reported that pesticides caused dermatitis (32.5%) in women exposed to OPs²¹. In consistent with our clinical manifestation's results, Hongsibsong et al.¹⁹, reported a significant health symptom, such as breathlessness, chest pain, dry throat, numbness, muscle weakness, cramping, dizziness, eye irritation, white/red rash, and white/red pimple among exposed farmers¹⁹.

The inhibition in activities of AChE and BuChE is commonly used as biomarkers for occupational and environmental exposure to OPs²². The advantages of these biomarkers for prediction of pesticide exposure may be attributed to their easy measurement, high sensitivity, and depending on the level of exposure²². We found a significant decrease in the levels of AChE and BuChE in

researchers occupationally exposed to OPs in comparison to the unexposed control group. Former studies have also reported decline in AChE and BuChE levels in pesticides exposed subjects relative to unexposed group^{23, 4}.

In addition to inhibitory effect of pesticides on cholinesterase activity, pesticides act as a powerful promoters of many cellular pathways. The long-lasting exposure to OPs leading to the generation of reactive oxygen species that exhaust and consume antioxidant agents present in the body^{24,25}. Disturbance of cellular oxidative balance related to pesticides exposure can be detected by measurement of lipid peroxidation by-product (MDA) and TAC level²⁶. In this study, there was a significant increase in MDA and a significant depletion in TAC between exposed researchers and controls **Table (1)**. The increase in MDA levels among pesticides exposed subjects was also reported in many former studies^{4, 27}. Other studies detected a significant decrease of TAC among exposed farmers to OPs^{28, 29}. In contrast, Ahmadi et al.³⁰, reported a significant increase in TAC in the pesticide-exposed group that could be due to the adaptive response to the generated free radicals³⁰.

In the present study, we detected the levels of inflammatory cytokines (IL-6, TNF- α and IFN- γ) that showed a significant increase in the exposed group **Table (1)**. Zaw et al.,³¹ reported that agricultural workers who were exposed to organophosphate pesticides have significantly higher serum levels of IL-6 in the exposed group than the non-exposed group³¹. On the other side, Taghavian et al.³² didn't find a significant increase of IL-6 among exposed group³². Massawe et al.,³³ reported an increase in levels of TNF- α , and IFN- γ in pesticides exposed group³³. Schaper et al.,³⁴ and Costa et al.³⁵ explained that OPs cause induction, production and the release of pro-inflammatory mediators, including cytokines such as TNF- α , interleukin (IL)-1, IL-6 and IL-10, or type 1 interferon (IFN)³⁵. The release of pro-inflammatory cytokines will lead to activation of immune cells and release of further cytokines (cytokine storm)³⁴. The Increased level of proinflammatory cytokines together with increased biomarkers of oxidative stress are indicators of immunotoxicity³⁵.

The present study indicated a positive correlation between MDA, IL6, and TNF levels and duration of pesticides exposure as illustrated in **Figure (2)**.

Our results reported a significant rise in MDA, IL6, and TNF levels with increasing duration of exposure. The BuChE level was significantly declined with more duration of exposure.

The correlation between measured biochemical biomarkers was represented in **Table (2)**. Regarding our results, there was a negative correlation between IL6 and both AChE and BuChE. Zaw³¹ et al., was in constant with our result that showing significant negative correlation between erythrocyte acetylcholinesterase activity and serum IL-6 level in pesticides-exposed group³¹. Our study indicated a significant positive correlation between MDA with IL-6 and TNF- α , while TAC was negatively correlated with TNF- α . **table (2)**. Mohammadzadeh. et al.³⁶ also reported significant increase in MDA level with IL-6 and TNF in rats treated by malathion³⁶. This correlation may be attributed to, depletion of antioxidant mechanisms that stimulate more production of inflammatory cytokines with increased risk of exposure hazards. In addition a positive correlation was noticed in our result between IL-6 and IFN- γ and between IL-6 and TNF- α **table (2)**. Cauvi et al.³⁷, has reported that in particular IFN- γ and IL-6 were required for the expression of each other in immunotoxicity³⁷. Both IL6 and TNF- α are important proinflammatory cytokines that are prominent in chronic inflammation³⁸.

From our study, Occupationally exposed researchers who complained from bronchial asthma was characterized with increased levels of MDA, IFN- γ , TNF- α **Table (3)**. Proskocil et al.³⁹, also were in line with us that OPs induce airway hyperactivity associated with increase of TNF- α ³⁹. Uysal and Karaman⁴⁰, reported that poisoning with OPs increases lipid peroxidation, disrupts the antioxidant system, induces histopathological changes, and increases apoptotic cell death in lung tissue⁴⁰. Zaw et al.³¹, found a significant negative correlation between serum IL-6 level and respiratory function parameters in OPs exposed group³¹.

According to our study, Neurological symptoms (muscular weakness and numbness) was associated with increased level of MDA, IL6, IFN- γ , TNF- α , and with decreased TAC level-**table (3)**. Our result agreed with Fareed et al.¹⁶, who reported a decreased level of TAC associated with muscular weakness in OPs exposed workers¹⁶. Caro-Gamboa et al²², reported that Malathion (organophosphorus

pesticides) can activate microglia to release inflammatory mediators (IL-6, TNF- α) and induce neuroinflammation²². Kulkarni et al.⁴¹, mentioned that the association of interferon-gamma with neurotoxicity⁴¹.

Exposed researchers with eye irritation have increased levels of MDA and IFN- γ **Table (3)**. General toxic effects of pesticides on the eyes are symptomatically classified as mainly external, resulting from direct ocular exposure. Subsequent absorption of OPs pesticides in the ocular tissue causes toxic effects, and/or internally through oxidative stress damage of ocular cells⁴². Ishikawa's epidemiological study revealed that 64 out of 7,435 (0.86%) farmers had impairment in ocular movements and abnormal contraction of the pupil in response to light, which improved after administration with antidotes of OPs (e.g atropine)⁴³.

In addition, we revealed exposed researchers with dermatitis related to significant rise in IL-6 **Table (3)**. Similarly, to the eye, OPs could affect skin mostly externally through direct contact and internally through disturbance in the immune system. Oxidative stress was recently involved in atopic dermatitis which was also accompanied by increased levels of proinflammatory cytokines⁴⁴.

Besides, incomplete or lack of use of personal protection equipment (PPE) by pesticides workers may potentiate the deleterious health effects in humans increasing the concentration of active chemicals in skin, mouth, eye, and respiratory tract, leading to an increase of pesticides absorption^{45,46}.

The use of PPE is considered the key to prevent adverse health symptoms of OPs exposure **Table (4)**. Our data suggested that OPs exposed researchers not constantly wearing a face masks during work showed a lower level of BuChE but higher levels of MDA, IL6, TNF- α than those regularly use face masks. Another study revealed a significant increase in adverse respiratory symptoms (shortness of breath and asthma) in OPs exposed workers not wearing respiratory PPE⁴⁷. We detected more BuChE inhibition level in exposed researchers not constantly wearing gloves than those regularly wearing gloves **Table (4)**. As the skin is an important route of absorption of OPs so, bare skin increases absorption which was detected by BuChE inhibition level in the exposed population. In addition, researchers not regularly wore an apron had an increased level of MDA as a

result of increased OPs exposure with subsequent increased oxidative stress. Hofmann et al.⁴⁸ reported that, not wearing chemical-resistant boots was strongly associated with BuChE inhibition level in OPs farmers⁴⁸.

Conclusion

From the current study can conclude that the severity of exposure to OPs was associated with consumption rate of antioxidant mechanisms and increased disturbance rate of inflammatory cytokines especially IL-6 and IFN- γ . There was a strong correlation between oxidative stress and alteration in cytokines levels within exposed researchers. This disturbance could trigger to a variety of immuno-disorder associated diseases. In this study bronchial asthma and neurological disorders was the most common clinical symptoms across exposed subjects. Furthermore, the use of PPE was observed to play an important in modulating immunotoxicity. So it's much recommended for researchers to maintain the required precautions and PPEs while handling with pesticides in their labs.

References

- 1- Espinoza Navarro O, Ponce La Rosa C, Bustos Obregón E., Organophosphorus pesticides: Their effects on biosentinel species and humans. Control and application in Chile. *Control and application in Chile. Int. J. Morphol.*, **35**(3),1069-1074 (2017)
- 2- Adekunle C. P. Akinbode S. , Akerele D.1 , Oyekale T. et al., Effects of Agriculture Pesticides utilization on Farmers health in Egeda local government area, Oyo State, Nigeria. *Nigerian Journal of Agricultural Economics (NJAE)*, **7**(1), 73-88 (2017).
- 3- Marina C. L. , Rui D. S. ,Nestor C. W. ,Guilherme R. et al., Developmental exposure to glyphosate-based herbicide and depressive-like behavior in adult offspring: Implication of glutamate excitotoxicity and oxidative stress *Toxicology* ,(**387**), 67-80 (2017).
- 4- Noshay MM, Saad-Hussein A, Shahy EM, El-Shorbagy HM, Taha MM, Abdel-Shafy EA. Assessment of anticholinesterase toxicity, oxidative stress and antioxidant status in carbamate and organophosphorus pesticides-exposed agricultural workers. *Toxicological & Environmental Chemistry*,**9**(3),205-9 (2017).
- 5- Singleton ST, Lein PJ, Dadson OA, McGarrigle BP, et al., Longitudinal assessment of

- occupational exposures to the organophosphorous insecticides chlorpyrifos and profenofos in Egyptian cotton field workers. *International journal of hygiene and environmental health*, **218**(2),203-11(2015).
- 6.Badr AM., Organophosphate toxicity: updates of malathion potential toxic effects in mammals and potential treatments. *Environmental Science and Pollution Research International*, 2020.
- 7- Cortés-Iza, Sandra Catalina, & Rodríguez, Alba Isabel. Oxidative stress and pesticide disease: a challenge for toxicology. *Revista de la Facultad de Medicina*, **66**(2), 261-267(2018).
- 8.Arab SA, Nikravesh MR, Jalali M, Fazel A., Evaluation of oxidative stress indices after exposure to malathion and protective effects of ascorbic acid in ovarian tissue of adult female rats. *Electronic physician*, **10**(5),6789 (2018).
9. Lakshmi Srinivasan ,Mary Catherine ,Harris Laurie and E. Kilpatrick, “ *Cytokines and Inflammatory Response in the Fetus and Neonate* “, Fetal and Neonatal Physiology , Vol.2 ,Chap.128,(Fifth Edition) , 1241-1254.e4 (2017)
- 10.Banks CN, Lein PJ., A review of experimental evidence linking neurotoxic organophosphorus compounds and inflammation. *Neurotoxicology*, **33**(3),575-84 (2012).
- 11- Damalas CA, Koutroubas SD., Farmers’ exposure to pesticides: toxicity types and ways of prevention. *Toxics.*, **4**(1), 1(2016).
- 12- Knedel M, Böttger R., Eine kinetische Methode zur Bestimmung der Aktivität der Pseudocholinesterase (Acylcholin-acylhydrolase 3.1. 1.8.). *Klinische Wochenschrift*, **45**(6),325-7(1967).
- 13.Ohkawa H., Ohishi N., Yagi K., Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical biochemistry*, **95**(2),351-81979.
- 14- Saad-Hussein A., Shafik Abdallah M., Shahy E.M., Fouda M.S., et al., Synergistic Oxidative Effects of Smoking and Pesticides Exposure on Reproductive Male Sex Hormones. *Journal of Biological Sciences*, **18** , 450-457(2018).
- 15- Sapbamrer R. and Nata S., Health symptoms related to pesticide exposure and agricultural tasks among rice farmers from northern Thailand. *Environmental Health and Preventive Medicine*, **19**(1),12-20 (2014).
- 16- Fareed M., Pathak MK., Bihari V., Kamal R., et al., Adverse Respiratory Health and Hematological Alterations among Agricultural Workers Occupationally Exposed to Organophosphate Pesticides: A Cross-Sectional Study in North India. *PLOS ONE*, **8**(8), 10.
- 17-Shaffo FC, Grodzki AC, Fryer AD, Lein PJ. Mechanisms of organophosphorus pesticide toxicity in the context of airway hyperreactivity and asthma. *Am J Physiol Lung Cell Mol Physiol.*, 2018 Oct 1;315(4):L485-L501
- 18- Drake MG, Lebold KM, Roth-Carter QR, Pincus AB, et al., Eosinophil and airway nerve interactions in asthma. *Journal of leukocyte biology*, **104**(1):61-7 (2018).
- 19-Hongsibsong S., Sittitoo N., Sapbamrer R., Association of health symptoms with low-level exposure to organophosphates, DNA damage, AChE activity, and occupational knowledge and practice among rice, corn, and double-crop farmers. *Journal of Occupational Health*, 16-0107 (2017).
- 20- Voorhees JR., Rohlman DS., Lein PJ., Pieper AA., Neurotoxicity in preclinical models of occupational exposure to organophosphorus compounds. *Frontiers in neuroscience*, **10**,590 (2017).
- 21- Farahat TM., Shaheen HM., Sanad ZF., Frag NA., Knowledge, attitudes, and practices of organophosphorus pesticide exposure among women affiliated to the Manshat Sultan Family Health Center (rural area) in Menoufia governorate: an intervention study. *Menoufia medical journal*, **29**(1), 115 (2016).
- 22- Caro-Gamboa LJ., Forero-Castro M., Dallos-Báez AE., Cholinesterase inhibition as a biomarker for the surveillance of the occupationally exposed population to organophosphate pesticides. *Ciencia y Tecnología Agropecuaria*, **21**(3),e1562(2020).
- 23.Mladenović M., Arsić BB., Stanković N., Mihović N., et al.,The targeted pesticides as acetylcholinesterase inhibitors: Comprehensive cross-organism molecular modelling studies performed to anticipate the pharmacology of harmfulness to humans in vitro. *Molecules*. **23**(9), 2192 (2018).
- 24- Mecdad AA, Ahmed MH, ElHalwagy ME, Afify MM., A study on oxidative stress biomarkers and immunomodulatory effects of pesticides in pesticide-sprayers. *Egyptian Journal of Forensic Sciences*, **1**(2), 93-8(2011).
- 25-Abdel-Shafy EA," Antioxidant status in pesticides-exposed agricultural workers", *Toxicology*, Chapter 5 , 43-51(2021) .

- 26-Gangemi S., Gofita E., Costa C., Teodoro M., et al., Occupational and environmental exposure to pesticides and cytokine pathways in chronic diseases. *International Journal of Molecular Medicine*, **38**(4), 1012-20 (2016).
- 27- Ntepe LJM, Habib R., Laure NJ., Raza S, Oxidative Stress and Analysis of Selected SNPs of ACHE(rs 2571598),BCHE (rs 3495),CAT (rs 7943316),SIRT1 (rs 10823108),GSTP1 (rs 1695), and Gene GSTM1,GSTT1 in Chronic Organophosphates Exposed Groups from Cameroon and Pakistan . *Int. J. Mol. Sci.*,**21**, 6432 (2020).
- 28- Hussein AS., Hafez SF., Shahy EM., Shaheen WA., et al., Prevalence of Neurological Abnormalities in Highly Educated Population Occupationally Exposed to Pesticides. *Int. J. Environ.*, **8** (2), 121-9 (2019).
- 29- Ghafouri-Khosrowshahi A, Ranjbar A, Mousavi L, Nili-Ahmadabadi H, Ghaffari F, Zeinvand-Lorestani H, Nili-Ahmadabadi A. Chronic exposure to organophosphate pesticides as an important challenge in promoting reproductive health: A comparative study. *J Edu Health Promot* ;**8**: 149 (2019)
- 30-Ahmadi N., Mandegary A., Jamshidzadeh A., Mohammadi-Sardoo M., et al., Hematological Abnormality, Oxidative Stress, and Genotoxicity Induction in the Greenhouse Pesticide Sprayers; Investigating the Role of NQO1 Gene Polymorphism. *Toxics.*, **6**(1):13 (2018).
- 31-Zaw T., Phyu MP., Kyaw S., Erythrocyte Acetylcholinesterase Enzyme Activity, Serum Interleukin-6 Level and Respiratory Function of Myanmar Agricultural Workers Exposed to Organophosphate Pesticides. *International Journal of Clinical and Experimental Physiology*, **7**(3), 107-11(2020).
- 32- Taghavian F., Vaezi G., Abdollahi M., Malekirad A.A., Comparative toxicological study between exposed and non-exposed farmers to organophosphorus pesticides. *Cell Journal (Yakhteh)*, **18**(1),89 (2016).
- 33-Massawe R., Drabo L., Whalen M., Effects of pentachlorophenol and dichlorodiphenyltrichloroethane on secretion of interferon gamma (IFN γ) and tumor necrosis factor alpha (TNF α) from human immune cells. *Toxicol Mech Methods.*, **27**(3), 223-235 (2017).
- 34- Schaper F. and Rose-John S., Interleukin-6: biology, signaling and strategies of blockade. *Cytokine & growth factor reviews*, **26**(5),475-87 (2015).
- 35- Costa C., Briguglio G., Catanoso R., Giambò F., et al. New perspectives on cytokine pathways modulation by pesticide exposure .*Current Opinion in Toxicology*, **19**, 99–104 (2020)
- 36-Mohammadzadeh, L., Hosseinzadeh, H., Abnous, K. et al. Neuroprotective potential of crocin against malathion-induced motor deficit and neurochemical alterations in rats. *Environ Sci Pollut Res* , **25**, 4904–4914 (2018).
- 37-Cauvi DM., Cauvi G., Toomey CB., Jacquinet E., et al., From the cover: interplay between IFN- γ and IL-6 impacts the inflammatory response and expression of interferon-regulated genes in environmental-induced autoimmunity. *Toxicological Sciences*, **158**(1),227-39 (2017).
- 38- Koelman L., Pivovarova-Ramich O., Pfeiffer A.F.H. et al. Cytokines for evaluation of chronic inflammatory status in ageing research: reliability and phenotypic characterisation. *Immun Ageing* **16**, 11 (2019).
- 39-Proskocil BJ., Grodzki AC., Jacoby DB., Lein PJ., Fryer AD., Organophosphorus pesticides induce cytokine release from differentiated human THP1 cells. *American journal of respiratory cell and molecular biology*, **61**(5),620-30 (2019).
- 40-Uysal M. and Karaman S., In vivo effects of intravenous lipid emulsion on lung tissue in an experimental model of acute malathion intoxication. *Toxicology and industrial health*, **34**(2),110-8 (2018).
- 41-Kulkarni A., Ganesan P., O'Donnell LA., Interferon Gamma: Influence on Neural Stem Cell Function in Neurodegenerative and Neuroinflammatory Disease. *Clin Med Insights Pathol*. **13**, 9-19 (2016).
- 42-Jaga K. and Dharmani C., Ocular toxicity from pesticide exposure: A recent review. *Environmental health and preventive medicine*, **11**(3):102-7(2006).
- 43-Ishikawa S., Ophthalmopathy due to environmental toxic substances especially intoxication by organophosphorus pesticides. *Nippon Ganka Gakkai Zasshi*, **100**(6),417-32 (1996).
- 44-Ji H. and Li XK., Oxidative stress in atopic dermatitis. *Oxidative medicine and cellular longevity*, **1**,(2016).
- 45-García-García CR., Parrón T., Requena M., Alarcón R., et al.,Occupational pesticide exposure and adverse health effects at the clinical, hematological and biochemical level. *Life sciences*, **145**,274-83 (2016).
- 46- Godoy F.R., Nunes H.F., Alves A.A., Carvalho

- W.F., et al., Increased DNA damage is not associated to polymorphisms in OGG1 DNA repair gene, CYP2E1 detoxification gene, and biochemical and hematological findings in soybeans farmers from Central Brazil. *Environmental Science and Pollution Research*, **26**(26),26553-62 (2019).
47. Muñoz-Quezada M.T., Lucero B., Iglesias V., Levy K., et al., Exposure to organophosphate (OP) pesticides and health conditions in agricultural and non-agricultural workers from Maule, Chile. *International journal of environmental health research*, **27**(1),82-93 (2017).
48. Hofmann J.N., Checkoway H., Borges O., Servin F., et al., Development of a computer-based survey instrument for organophosphate and N-methyl-carbamate exposure assessment among agricultural pesticide handlers. *Annals of occupational hygiene*, **54**(6),640-50 (2010).