



Full Profiling of the Orange By-Products and Their Potential Effects with Chromium on Obesity Complications in Rats



CrossMark

Ebtehal A Altamim*

Associate Professor of Nutrition & Food Science, Physical Sport Science Department, Education College,
Princess Nourah Bint Abdulrahman University, Riyadh, Kingdom of Saudi Arabia

Abstract

A disorder identified as obesity includes having too much body fat. It is a medical ailment that increases the risk of several syndromes and other health issues. Around the world, 19.5% of the adult population is obese, which is a widespread condition. The current study seeks to explore the chemical constituents, bioactive substances, antioxidant activity, and impacts on obesity problems in rats of orange by-products, peel, and pulp integration with chromium. While orange pulp has larger amounts of carbohydrates, minerals (Mg, Fe, and Zn), vitamins (E, B1, and B6), and bioactive substances than orange peel, the latter has higher levels of total protein, fiber, minerals (Na, K, Ca, and P), vitamins (A and C), total polyphenols, flavonoids, and carotenoids. In contrast to the control normal group, rats fed a diet-stimulated obesity diet (DIO) had a higher body weight (416.76%), blood leptin (142.84%), glucose (95.17%), and hyperlipidemia (cholesterol (219.77%), triglycerides (131.22%), and low-density lipoprotein cholesterol (341.59%). Because orange peel and pulp contain antihyperlipidemic and anti-diabetic substances that lower serum total cholesterol, triglycerides, and low-density lipoprotein cholesterol and raise high-density lipoprotein cholesterol, consuming them alone or in combination with chromium partially reversed these negative changes and decreased the likelihood of atherogenesis. When orange peels and pulp were combined with chromium, there showed noticeable synergistic effects. These results lay a foundation for the use of food industry by-products, with or without chromium, in the early detection and treatment of obesity. Additionally, the current study suggests adding orange by-product powder and/or extracts to our everyday meals, beverages, dietary supplements, and pharmaceutical formulations.

Keywords: Orange peel & pulp; Composition; Serum lipid profile; Atherogenic index; Leptin; Insulin.

1. Introduction

A metabolic disease known as obesity is brought on by an excess of calories consumed and a lack of physical exercise [1]. Obesity is associated with several chronic circumstances, containing hypertension, dyslipidemia, and coronary heart ailments [2,3]. Body mass index (BMI), which is expressed as the weight in Kgs divided by the square of the height in meters (kg/m²), is a standard tool for determining the prevalence of bulky and obesity. Corpulence is described as a BMI of over 25 kg/m², and obesity as a BMI of over 30 kg/m². These indicators serve as common reference points for evaluation; however, illness risks in all groups might gradually rise with decreasing BMI levels [4,5]. Over half of adults in developing countries are overweight,

and 19.5% of adult populations are obese, according to the Organization for Economic Co-operation and Development member countries (OECD) [6]. Additionally, according to the World Health Organization (WHO), there are more than a billion overweight individuals worldwide [7]. At least 115 M of them are from developing nations, and there are at least 300 M of them who are clinically obese. Additionally, statistical data shows that the obesity dilemma has grown from 12–20% for males and from 16–25% for females [8]. According to several studies, obesity affects around 15–20% of middle-aged people in Europe [9], while it affects up to 40% of people in the USA as 3,00,000 untimely fatalities each year [10]. Patients who are obese have a higher risk of morbidity and death than those who have an appropriate body weight [11]. Significant improvements in a variety of co-morbid illnesses are linked to even moderate

*Corresponding author e-mail: dr.ebaltamim@gmail.com; (Ebtehal A Altamim)

Receive Date: 22 November 2022, Revise Date: 03 January 2023, Accept Date: 08 January 2023.

DOI: [10.21608/ejchem.2023.176415.7221](https://doi.org/10.21608/ejchem.2023.176415.7221)

©2023 National Information and Documentation Center (NIDOC)

weight loss in the scale of 5–10% of the starting body weight [12].

Accordingly, seeking effective methods to prevent and/or treat obesity and its associated metabolic diseases is highly demanded. Meanwhile, many pharmaceuticals have been used in this direction, but unfortunately lots of them were not effective, alongside their harmful side effects and the excessive cost. Therefore, valorizing the natural sources to be used as a safe alternative to those substances was recently spotlighted. Amongst, the plant parts, especially, food processing by-products are the commonly and most important sources to be used and most [13,14,15].

According to estimates, 18.14 M tons of wastes are annually produced because of industrialized agriculture in the Arab world, with leftovers from the production of fruits and vegetables accounting for around 6.14% of this total. Fruits and vegetables that have been processed produce significant volumes of waste products, including peels, seeds, stones, meals, etc. It is commonly recognized that agricultural by-products are high in dietary fibre, and some of them also include significant amounts of coloring agents, antioxidants, or other chemicals that have beneficial effects on health [16]. Citrus and oranges, two of the most popular fruits, are a significant source of food by-products.

The citrus fruit known as orange (*Citrus sinensis* L.) is a member of the *Rutaceae* family and an evergreen declining tree. In tropical and subtropical climates, orange trees are frequently grown for their fruit, which is skinned or chopped (to prevent the bitter rind) and consumed entire, or treated to obtain orange juice, as well as their aromatic peels. Around the world, 68.5 M tons of oranges were cultivated in 2008, mostly in Brazil and the US state of Florida. Oranges were grown in China by 2500 BC and came from Southeast Asia [17]. Orange fruits are entirely peeled and utilized for food, orange juice production, or fragrance production. It includes a variety of vital vitamins, minerals, and nutrients that support healthy growth and development. Peels, seeds, and pulp are the waste products of the citrus processing business. Orange peels comprise around 50% of the weight of the fresh fruit and are high in pectin, cellulose, and hemicellulose but low in protein (5.8%) [18]. 50–70% of the fresh weight of the initial fruit is invented of the pulp. The peel (60–65%), interior tissues (30–35%), and seeds (0.5–10%) are all parts of it. The essential oils from the peel are abundant in bioactive substances such coumarins, flavonoids, carotenes, terpenes, and

linalool [19]. Natural phenolic chemicals, especially flavanone glycosides, principally naringin, hesperidine, narirutin, and neohesperidine, are abundant in it, making it stand out as a particularly rich source [20]. Citrus flavonoids have been linked to a broad range of biological functions, involving antibacterial, anticancer, antidiabetic, and antioxidant actions [21]. Additionally, Lv et al. [22] looked at how orange-peel essential oil affected acute otitis media rats' oxidative stress. The blood levels of malondialdehyde (MDA), immunoglobulin A (IgA), immunoglobulin G (IgG), and immunoglobulin M (IgM) could all be decreased, and antioxidant enzyme activity could be increased by using orange peel essential oil. For the above reasons, this study was performed to examine the possible effects of chromium consumption together with orange byproducts (peels and pulps) on the complications of obesity in rats. The scope of this study will also include the chemical composition, minerals, vitamins, and bioactive substances of such plant sections.

2. MATERIALS AND METHODS

2.1. Materials

2.1.1. Orange fruits

During the 2020 harvesting season, orange (*Citrus sinensis* L.) fruits were purchased from Saudi Arabian regional marketplaces. The gathered samples were taken to the lab and utilized right away to prepare the peels and pulps.

2.1.2. Chemicals

The casein was collected from Morgan Chemical Company. The chromium picolinate (CHROMIUM® capsules) were purchased from EPACO Company. The typical bioactive chemicals (gallic acid (GA), Butylated hydroxytoluene (BHT), Cu_2O_4 , β -carotene, and α -tocopherol) and vitamins (B₁, B₂, B₆, C, and E) were purchased from Sigma Chemical Co., St. Louis, MO. All other chemicals and solvents were obtained from analytical or HPLC-rating suppliers (Fisher, UK). The mobile parts, reagent solutions, and standards were all made with dH_2O (Milli-Q 18.2 M).

2.1.3. Machines

Throughout this investigation, the UV-160A from Shimadzu Corporation in Kyoto, Japan, was used to detect absorbance for several assays. Additionally, a Perkin-Elmer Model 2380 atomic absorption spectrophotometer from Waltham, Massachusetts, USA, was utilized to determine the minerals. Additionally, a Consta Metvic 4100 pump, a Spectra Series AS100, a Spectra System UV 1000 UV/Visible Spectrophotometer Detector, a Spectra System FL 3000, and a PC 1000 system software were utilized with SP Thermo Separation Products Liquid Chromatography (Thermo Separation Products, San Jose, CA, USA). Adsorbosil C₁₈ (5 µM, 100 mm × 4.6 mm I.d.) for the analysis of water-soluble vitamins and regular Ultrasphere Si (5 µM, 250 mm × 4.6 mm I.d.) for the analysis of fat-soluble vitamins were the columns that were employed (Alltech, Deerfield, IL, USA).

2.2. Methods

2.2.1. Preparation orange peels and pulps

Orange fruits were manually cleaned, peeled, and pressed to get pulp from the orange flesh. In two phases, at 50 °C for 4 h and then 40 °C for 6 h, the collected peels and pulps were dehydrated in a warm air oven (Horizontal Forced Air Drier, Proctor and Schwartz Inc., Philadelphia, PA). The dehydrated peels and pulps were processed at a high blender rapidly to create a fine powder. The items kept for usage must pass through an 80-mesh filter.

2.2.2. Preparation of orange peel & pulp extracts

For their many forms of extracts, orange peel & pulp powder were utilized as follows: A beaker containing 20 g of dried orange peels or pulps and 180 mL of EtOH (80%, v/v) was homogenized before being shaken at 200 rpm for 1 h at RT (Unimax 1010, Heidolph Instruments GmbH & Co. KG, Germany). Whatman No. 1 filter paper was used to separate the extract from the residue during filtration. The left-over residue was extricated twice more isolation, and the two extracts were gathered after that. The remaining solvent was isolated with the aid of a rotary evaporator at 45 °C and lower pressure (Laborata 4000; Heidolph Instruments GmbH & Co. KG, Germany).

2.2.3. Chemical analysis of orange peel and pulp powders

Using the techniques outlined in the AOAC, orange peel & pulp samples were examined for proximate chemical composition, containing moisture, protein (T. N. 6.25, micro-kjeldahl assay *via* autoloading device, Velp company, Italy), fat (Soxhlet miautomatic system Velp company, Italy), ash, fibre, and dietary fibre contents [23]. Difference-based calculations of carbohydrates was of:

$$\text{Carbohydrates (\%)} = 100 - (\% \text{moisture} + \% \text{protein} + \% \text{fat} + \% \text{Ash} + \% \text{fiber})$$

2.2.4. Determination of minerals content

Using the method explained by Singh *et al.* [24] the minerals content of orange peel and pulp powder samples was tested as follows: The Kjeldahl digestion unit received 0.5 g of the defatted material to be processed, and each tube received 6 mL of the tri-acid combination (HNO₃, HClO₄, and H₂SO₄ acid in the ratios of 20: 4: 1 v/v, orderly). The tubes' innards were slowly absorbed as observed: for 30 min / 70 °C, for 30 min / 180 °C, then for 30 min / 220 °C. The combination was digested, chilled, and then dispersed in dH₂O. The quantity was then raised to 50 mL in a volumetric beaker. A selection of minerals was examined in aliquots following filtering *via* ashless filter paper with the aid of atomic absorption spectrophotometer.

2.2.5. Vitamins determination

Water soluble vitamins (B and C) were extracted from orange peel and pulp powders in accordance with Moeslinger *et al.* and Epler *et al.* procedures [25,26], while fat soluble vitamin (E) was done so in accordance with Epler *et al.* [26] and Hung *et al.* methods [26,27]. Vitamins C, B₁, B₂, B₆, and E recoveries were 89.98±3.76, 85.67±2.98, 88.65±2.09, 84.99±1.83, and 90.03±1.21%, separately under the chromatographic conditions utilized in those procedures.

2.2.6. Bioactive compounds determination

Singleton, Rossiand and Wolfe *et al.* used the Folin-Ciocalteu reagent to evaluate the total phenolics in orange peel and pulp powder samples [28,29]. Results are presented as equivalents and gallic acid (GAE). Utilizing the assay described by Lichtenthaler [30], the sum carotenoids in the 80% acetone extract were calculated and expressed as g of carotenoid/g of dry extract. Utilizing the colorimetric technique established by Zhishen *et al.* [31], the value of total flavonoids was calculated and represented as mg of catechin equivalent, CAE, per g of dry extract as follow:

$$Y = 0.0003x - 0.0117, r^2 = 0.9827, \text{mg of CA/g of dehydrated extract}$$

2.2.7. Antioxidant activity (AA)

A modified version of the process Marco outlined was used to assess the AA of orange peel and pulp extracts and standards (α-tocopherol and BHT) [32]. Using the Al-Saikhan *et al.* equation [33], antioxidant activity (AA) was all measured as a percentage inhibition compared to control as follows:

$$AA = (R_{\text{control}} - R_{\text{sample}}) / R_{\text{control}} \times 100$$

R_{control} and R_{sample} were the levels of β -carotene bleaching in the reactant blend in the absence of antioxidant and with byproducts' extract, orderly.

2.3. Biological experiments

2.3.2. Animals

Animals utilized in this study, adult male albino rats (100 ± 10 g per each) were collected from Animal Lab, King Saud University, Riyadh, Saudi Arabia.

2.3.3. Basal Diet

According to AIN, the basic diet should consist of the following ingredients: Protein (10%), corn oil (10%), vitamin mixture (1%), mineral combination (4%), choline chloride (0.2%), methionine (0.3%), cellulose (5%), and the remainder should be corn starch (69.5%). According to AIN, the employed vitamin and metals mixed compounds was created [34].

2.3.4. Experimental design

The National Research Council, Commission on Life Sciences, and Institute of Laboratory Animal Resources regulations were followed in all biological research [35]. Rats ($n=48$ rats) were kept in separate rope cages in a room that was kept at a constant temperature of 25 ± 2 °C and at a damp level of 50–60%. Prior to the trial, all rats were given a baseline diet for one week to help them become used to it. The rats were separated into two major groups after a one-week period, with the initial group (Group 1, 6 rats) continuing to consume a baseline diet while the other major group (42 rats) was fed a diet-induced obesity (DIO) diet (Product No. D1245, Research Diets, Inc. NJ, Protein, 24 g/100 g (Kcal, 20%) carbohydrates, 41 g/100 g (Kcal, 35%) and fat, 24g/100 g (Kcal, 45%) for 4 weeks which categorized into 7 sub-groups as follow: group (2), which received DIO as a positive control; group (3), which received DIO plus Chromium Picolinate 80 g/kg/d by stomach tube; group (4), which received DIO containing 15% orange peels powder (OPP); group (5), which received DIO containing 15% orange pulp powder (OPuP); group (6), which received DIO containing mixture of OPP plus chromium; group (7); and group (8), which received DIO containing mixture. Rats were weighed before the trial began, then every week, and lastly at the end of the experiment.

2.3.5. Biological assessment

Throughout the trial time, daily dietary records and weekly body weight records were kept (8 weeks). According to Chapman *et al.* [36], the body weight gain (BWG, %), food intake (FI), and food

efficacy ratio (FER) were calculated by the subsequent equations:

$$\text{BWG (\%)} = (\text{Ending weight} - \text{Preliminary weight}) / \text{Preliminary weight} \times 100$$

$$\text{FER} = \text{g that increase in body weight (g/28 d)} / \text{g that feed consumption (g/28 d)}$$

2.3.6. Blood testing

The abdominal aorta was used to collect blood samples after 12 h of fasting at the conclusion of the 28-d trial, and rats were scarified while being sedated with ether. Blood samples were put in clean, dry separator tubes and allowed to clot at RT before being spun for 10 min at 3000 rpm to split up the serum, using the Drury and Wallington [37]. The serum was meticulously extracted, moved to sterile, clean tubes, and then frozen at -20 °C pending analysis.

2.3.7. Hematological analysis

Hematological analysis involving serum total cholesterol (CHO), triglycerides (TGs), high-density lipoprotein cholesterol (HDL-C) and total lipids (T. lipids); blood leptin, insulin, and glucose; and liver cholesterol, TGs, T. lipids and glycogen using the methods of Everds and, Ramaiah [38]. According to Lee and Nieman [39], the following formulas were utilized to multiply very-low-density lipoprotein cholesterol (VLDL-C), low-density lipoprotein cholesterol (LDL-c), and phospholipids as follow: $\text{VLDL-C} = \text{TG} / 5$, $\text{LDL-C} = \text{CHO} - \text{HDL-C} - \text{VLDL-C}$, and phospholipids = T. lipids - (TGs – CHO). Also, the atherogenic index was determined using the formula provided by Castelli and Levitar [40]: $\text{Atherogenic index} = (\text{CHO} / \text{HDL-C and LDL-c} / \text{HDL-C}$

2.4. Statistical analysis

Using the Student *t*-test and the statistical software MINITAB 12, all data were statistically evaluated (Minitab Inc., State College, PA). The data was shown as means with a standard deviation (SD). $P \leq 0.05$ was utilized to verify if differences between treatments were significant [41].

3. RESULTS AND DISCUSSION

3.1. Proximate chemical composition of orange by-products

The approximate composition of orange by-products was tabulated in **Table 1**. It could be noted that the peel contained higher values of total protein and fiber, while the pulp had the higher values of carbohydrates and moisture. Both contained similar amounts of ash and fat. These data are partially matched with that reported by Abd-Elwahed [16].

Related results have been reported in different studies, while many factors affect the contents of orange's peel and pulp like climate, citrus variety, agricultural, and analytical methods [42]. Owing to their extra nutritional value, rich resources of protein, fibre, and ash, and data from this study and others, these plant components might be employed successfully in food technology and/or nutritional applications. It is also a low-fat calorie food, subsequently more suitable for humans in diet and cardiovascular diseases. Additionally, orange by-products are characterized by a high fiber content. Although fiber is indigestible, it plays a significant nutritional role since it helps to provide bulk to stool and aid in the movement through the digestive tract.

Table 1. Approximate composition of orange by-products (on dry weight, DW).

Parameters (g/100g)	OPP	OPuP
Ash	5.18 ± 0.35	5.05 ± 0.51
Total fats	2.28 ± 0.21	2.18 ± 0.32
Total proteins	11.58 ± 1.56	7.63 ± 0.77
Total carbohydrates	57.73 ± 3.25	59.17 ± 2.21
Crude fiber	12.66 ± 0.65	9.29 ± 0.74
Total moisture	10.57 ± 1.32	16.68 ± 1.67

Each value signifies the Mean of three replicates ± SD. Means under the same row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder.

3.2. Minerals' concentration of orange by-products

Data of mineral contents of orange by-products has been illustrated in **Table 2**. Data indicated that the peel contained higher values of Na, Ca, and P, while the pulp had higher values of K, Mg, Fe, and Zn. Minerals such as K, Ca, P, and Mg are presented to be the major elements as they found in high concentrations of orange by-products. However, Na is less in orange by-products; thus, orange by-products could be recommended as a useful source for patients with hypertension. Trace metals are also found in orange by-products in considerable amounts in their inorganic form. All those trace elements are biologically very vital to the human body through prevention and/or fighting many diseases including anemia, immunodeficiency cancer, and atherosclerosis [43]. Zn's function in cell development, division, and maturation, cell membrane stability, as well as DNA and RNA production, all contribute to its significance in the immune system [44,45]. The primary function of Fe, which is a component of hemoglobin in red blood cells, is the delivery of O_2 from the lungs to the tissues of all bodily organs. Additionally, it is vital for DNA synthesis and the functioning of the human immune system [45].

Table 2. Minerals' concentration of orange by-products (on dry weight, DW).

Minerals (mg/100 g)	OPP	OPuP
Sodium	3.09 ± 0.12	2.66 ± 0.20
Potassium	212.89 ± 14.10	131.66 ± 9.16
Calcium	161.40 ± 10.65	109.05 ± 7.57
Phosphorous	21.94 ± 1.67	19.80 ± 2.02
Magnesium	22.88 ± 1.03	42.99 ± 3.27
Iron	108.65 ± 9.68	175.88 ± 14.29
Zinc	47.25 ± 3.09	90.43 ± 6.10

Each value signifies the Mean of three replicates ± SD. Means under the same row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder.

3.3. Vitamins content of orange by-products

The vitamin content of orange by-products is displayed in **Table (3)**. The peels had the highest concentrations of vitamins C and β -carotene, whereas the pulp held the highest concentrations of vitamins E, B₁, and B₆. From a nutritional perspective, vitamins are necessary for life since they support development and good health. Vitamin B₁, B₂, and B₆ are all found in significant amounts in orange byproducts. In terms of structure and function, vitamins B constitutes a varied group. It has a role in the metabolism of several amino acids, including histidine, serine, methionine, and glycine. On the other hand, there are the anti-inflammatory minerals, such β -carotene (found in orange byproducts) and vitamins C, E, and Nutritional Reference Values (DRVs).

Table 3. Vitamins content of orange by-products (on dry weight, DW).

Vitamins	OPuP	OPP
Vitamin A (β -carotene, $\mu\text{g}/100\text{g}$)	292.55 ± 17.56	180.79 ± 10.21
Vitamin C (mg/100g)	39.18 ± 3.76	23.38 ± 2.42
Vitamin E ($\mu\text{g}/100\text{g}$)	1.72 ± 0.23	5.71 ± 0.53
Vitamin B ₁ (mg/100g)	0.29 ± 0.05	0.42 ± 0.07
Vitamin B ₂ (mg/100g)	0.10 ± 0.01	0.15 ± 0.04
Vitamin B ₆ (mg/100g)	0.264 ± 0.031	0.360 ± 0.028

Each value signifies the Mean of three replicates ± SD. Means under the similar row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder.

3.4. Bioactive compounds of orange by-products

Bioactive compounds of orange by-products were illustrated in **Table (4)**. Data noticed that the total polyphenols, flavonoids, and carotenoids were higher in the pulp. Our data partially agreed with that noted by Abd-Elwahed [16]. Numerous of our earlier

studies, as well as other research, have demonstrated the critical role those bioactive substances, such as the phenolics, flavonoids, and carotenoids found in orange by-products in this study, play in averting and/or treating several ailments, including diabetes, atherosclerosis, cancer, obesity, bone, and ageing [14, 46]. The miraculous antioxidant properties of these chemicals are primarily responsible for all the benefits. Due to their rich supply of both nutritional and non-nutritional substances, all these elements combined to make orange by-products a full bundle of nutritious food.

Table 4. Bioactive compounds of orange by-products (on dry weight, DW).

Parameters (mg/100g)	OPP	OPuP
Total polyphenols	0.37 ± 0.09	0.44 ± 0.11
Flavonoids	0.54 ± 0.16	0.91 ± 0.15
Carotenoids	0.274 ± 0.03	0.389 ± 0.04

Each value signifies the Mean of three replicates ± SD. Means under the similar row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder.

3.5. Antioxidant activity (AA) of orange by-products

Table (5) displays the antioxidant activity (AA) of orange by-products. It was evident from such data that changes in the orange by-products ranged from 73.65-

80.67%, which was different ($p \leq 0.05$). Due to the abundance of phytochemicals and nutrients in all the examined sections, they all showed robust activity (phenolics, flavonoids, carotenoids, essential oils, vitamins, and other components). These outcomes are constant with those described by Abd-Elwahed [16], who discovered that various by-products of food processing, such as orange peels and others, have a strong antioxidant activity because of their high presence of phenolic constituents. In a related study, Velioglu *et al.* revealed that there was a statistically significant association between the total phenolic content and antioxidative activity of twenty-eight plant products, including plant by-products [47]. Additionally, several investigations revealed a favorable and substantial ($p \leq 0.01$) correlation between total phenolics and AA in various plant sections [46,48,49,50]. Antioxidants aid in defending cells against "oxidative stress," a physiological condition that may be harmful (injury to healthful cells or DNA by peculiar electrons recognized as free radicals). Chronic disorders such as cancer, heart syndrome, diabetes, rheumatoid arthritis, obesity, and age-related ailments such as neurodegenerative diseases like Parkinson's and Alzheimer's are influenced by oxidative stress [13,15,51,52,53,54]. The orange by-products were advised to be employed successfully in several medicinal applications for all of these and other reasons.

Table 5. Antioxidant activity (AA) of orange by-products and standards.

Parameters	OPP	OPuP
AA (%)	73.56 ± 4.52	80.67 ± 3.09
AA [% of BHT (50 mg/mL)]	83.01 ± 2.31	91.03 ± 0.99
AA [% of BHT (100 mg/mL)]	87.28 ± 1.45	84.81 ± 2.03
AA [% of α -tocopherol (50 mg/mL)]	75.05 ± 2.91	82.30 ± 1.79
BHT (50 mg/mL)	88.62 ± 0.49	88.62 ± 0.49
BHT (100 mg/mL)	95.11 ± 0.81	95.11 ± 0.81
α -tocopherol (50 mg/mL)	98.02 ± 0.57	98.02 ± 0.57

Each value signifies the mean of ten replicates ± SD. Mean values with the different superscript letters in the similar row mean significantly different at level $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder; BHT, butylated hydroxytoluene.

3.6. Biological studies

3.6.1. Effect of orange by-products on BWG, FI, and FER of obese experimental rat groups

Data in **Table (6)** showed the BWG, FI, and FER of the obese investigational rat groups. The values of these parameters had increased in the positive control obese rat group consuming the DIO, while they had significantly ($p \leq 0.05$) decreased in the chromium, orange peel, orange pulp, orange peels with chromium, orange pulps with chromium, and mixed orange peels and pulps with chromium groups. The mixed orange peel and pulp with chromium group listed as the lowest values compared to the positive control group.

This study demonstrates obesity-reducing effects of the dietary parameters of chromium and both orange peels and pulps. Chromium is vital for maintaining the ordinary carbohydrate and lipid metabolism and attenuating certain DIO impacts, primarily body fat accretion. Orange by-products contain considerable amounts of bioactive compounds (phenolics, carotenoids, flavonoids, etc.) and vitamins (A, C, and E) that have strong anti-inflammatory and antioxidant features and exhibited the positive effects of such plant parts in regarding the control of the obesity [14,15,16,55,56]. Orange peel and pulp also contain high amount of fiber which positively affects the regulation of intestinal capacity, disease prevention

and as a therapeutic factor in diverse diseases, especially the metabolic disorders [57,58].

Table 6. BWG, FI, and FER of obese investigational rat groups.

Parameters	Negative control	Positive control	Chromium	OPP	OPuP	OPP + chromium	OPuP + chromium	OPP+ OPuP+ chromium
BWG	34.67±1.12e	179.16±4.61a	104.55±3.33c	110.33±3.66b	113.07±3.11b	49.66±2.01d	45.44±2.11d	36.99±2.05e
FI	22.98±2.98c	39.05±3.41a	30.33±3.05b	31.01±3.10b	31.51±3.33b	24.44±3.21c	24.60±3.71c	23.59±3.61c
FER	0.024±0.003d	0.085±0.002a	0.057±0.01b	0.059±0.001b	0.060±0.003b	0.033±0.004c	0.30±0.001c	0.026±0.002cd

Each value represents the Mean ± SD (n=6). Means under the same row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder, BWG, body weight gain; FI, feed intake; FER, feed efficacy rate.

3.6.2. Impact of orange by-products on serum lipid profile of obese investigational rat groups

Data in **Table (7)** demonstrated how orange byproducts affected the groups of obese experimental rats' blood lipid profiles. The positive control obese rat group that ingested the DIO exhibited high values for the blood lipid profile (CHO, TGs, LDL-c, and VLDL-c) and low HDL-c, according to the current findings. Chromium, orange peels, and orange pulps treatment decreased their prominent levels and elevated HDL-c. When related to the positive control group, the administration of chromium with orange peels or pulps combination obviously had synergistic effects on lowering CHO and LDL-c levels. Citrus flavonoids

may have a lipid-lowering effect as because of their antioxidant properties, suppression of the absorption of oxidized LDL by macrophages, and decreased LDL aggregation and oxidation of LDL cholesterol. When compared to a hypercholesterolemic diet, diets supplemented with 10 and 20% orange powder significantly lowered T. lipids, CHO, TGs, and LDL-c in rats [59]. In the same context, modelling based on systematic reviews of RCTs reveals that modest and sustained weight loss (5-10 kg) in individuals with overweight or obesity is linked to decreased levels of LDL-c, CHO, and TGs as well as elevated levels of HDL-c [60,61,62].

Table 7. Serum lipid profile of the obese experimental rat groups.

Parameters (mg/dL)	Negative control	Positive control	Chromium	OPP	OPuP	OPP +chromium	OPuP + chromium	OPP+ OPuP + chromium
CHO	123.56±6.78e	395.11±5.77a	197.55±4.21b	191.77±3.99b	190.55±3.61b	170.81±4.11c	165.33±5.11c	155.11±5.31cd
TGs	87.12±3.02c	201.44±7.33a	115.40±6.11b	110.44±5.19b	112.20±5.31b	105.22±5.23b	103.77±5.11bc	95.16±4.01c
HDL-c	30.67±2.45a	21.11±1.77b	38.99±2.41a	38.01±2.99a	37.81±2.81a	40.31±3.01a	40.71±3.66a	34.21±3.25a
LDL-c	75.47±3.15e	333.72±7.96a	135.48±4.21b	131.67±4.77b	13.31±4.31b	109.46±5.1c	103.87±5.82c	92.87±4.71cd
VLDL-c	17.42±0.99c	40.28±2.11a	32.08±1.45b	22.9±1.66b	22.44±1.34b	21.04±1.77b	20.75±1.36bc	19.03±1.22c

Each value represents the Mean ± SD (n=6). Means under the same row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder; CHO, cholesterol; TGs, triglycerides; HDL-c, high density lipoprotein cholesterol; LDL-c, low density lipoprotein cholesterol; VLDL-c, very low-density lipoprotein cholesterol

On the other side, data in **Table (8)** illustrated that the increase of atherogenic indexes (CHO/HDL-c and LDL-c/HDL-c) of positive control obese rat group, while administration of chromium, orange peels & pulps or a combination of orange peels and/or pulps with chromium clearly showed their synergistic effects to reduce CHO and LDL-c and raised the HDL-c. Additionally, there was an increase in serum T. lipids and lower phospholipids in the positive control obese rat group, while groups administrated with chromium, orange peels, orange pulps or a combination of the two with chromium reduced T.

lipid levels (**Table 9**). The chromium, orange peels and orange pulps groups showed lower values of phospholipids, while the orange pulps with chromium group and the orange peels and pulps with chromium group showed higher values of phospholipids compared to the positive control group. Several studies on diabetes illustrated that supplementation with chromium ion (III) reduced blood glucose concentrations, CHO, LDL-c, TGs, and non-esterified fatty acids levels; as well as enhanced the concentrations of HDL-c [63].

Table 8. Atherogenic parameters (CHO/HDL-c and LDL-c/HDL-c) of the obese experimental rat groups.

Parameters	Negative control	Positive control	Chromium	OPP	OPuP	OPP + chromium	OPuP + chromium	OPP+ OPuP+ chromium
CHO/HDL-c	4.03±0.32c	18.71±1.55a	5.6±0.41b	5.04±0.53b	5.03±0.46b	4.23±0.35bc	4.06±0.45bc	3.58±0.44c
LDL-c/HDL-c	2.46±0.20c	15.80±1.60a	3.47±.31b	3.46±.37b	3.44±0.35b	2.71±0.22b	2.55±0.28bc	2.14±0.26c

Each value represents the Mean ± SD (n=6). Means under the same row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder.

Table 9. Serum total lipids and phospholipids of the obese experimental rat groups.

Parameters (mg/dL)	Negative control	Positive control	Chromium	OPP	OPuP	OPP + chromium	OPuP + chromium	OPP+ OPuP+ chromium
T. lipids	289.67±11.39c	703.93±33.61a	392.11±11.77b	387.41±13.41b	379.75±14.55b	380.71±19.33b	391.79±17.77b	374.55±16.44b
Phospholipids	78.99±9.42b	107.38±9.66b	79.16±7.22d	85.21±8.75c	77.01±7.42d	104.68±10.21b	122.69±13.61a	124.28±14.25a

Each value represents the Mean ± SD (n=6). Means under the same row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder; T. lipids, total lipids.

3.6.3. Effect of orange by-products on blood leptin, insulin, and glucose levels of obese investigational rat groups

According to data in **Table (10)** the positive control group of obese rats had significantly higher ($p \leq 0.05$) blood levels of leptin and glucose and lower insulin values, whereas the treatment of chromium, orange peels, and orange pulps boosted leptin and glucose

levels and decreased insulin levels. In the orange peels with chromium, orange pulps with chrome, and orange peels and pulps with chromium groups related to the positive control group, there was a decrease in leptin and glucose and an increase in insulin. Studies have revealed that obese people have greater serum leptin levels, and that chromium can lower blood sugar and cholesterol by increasing glucose and insulin tolerance [64].

Table 10. Blood leptin, insulin, and glucose levels of the obese investigational rat groups

Parameters	Negative control	Positive control	Chromium	OPP	OPuP	OPP + chromium	OPuP + chromium	OPP+ OPuP+ chromium
Leptin (ng/l)	46.45±2.92d	112.80±6.33a	63.14±2.77b	60.79±3.41b	61.11±3.22b	55.17±2.96c	56.37±3.01c	54.11±3.19c
Insulin (mg/dL)	16.41±3.08a	10.11±1.22d	12.80±1.11c	12.59±1.61c	13.55±1.33ac	14.11±1.41b	14.03±1.92b	14.77±1.80b
Glucose (mg/dL)	110.56±4.09e	215.78±5.33a	160.71±4.55b	168.40±5.76b	163.33±6.01b	140.76±4.99c	145.84±5.12c	130.77±3.98d

Each value represents the Mean ± SD (n=6). Means under the same row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder.

3.6.4. Effect of orange by-products on liver cholesterol, total lipids, triglycerides (TGs), and glycogen of the obese investigational rat groups

The positive control obese rat group revealed significant changes ($p \leq 0.05$) in increasing liver cholesterol and T. lipids and decreasing TG and glycogen values, while all other groups showed significant changes in decreasing liver cholesterol and T. lipids and increasing glycogen, according to data in **Table (11)**. When equated to the positive control group, TGs were enhanced when orange peels and pulps were administered together with chromium. Considering this, hesperidin, a flavonoid that aids in

lowering cholesterol and TGs, is present in orange peels and pulps [65]. Additionally, oranges' ability to decrease cholesterol is due to the limonene component in their peels. Furthermore, polymethoxylated flavones found in citrus fruit peels can lower cholesterol more effectively and safely than certain prescription drugs. On the other hand, by lowering the levels of T. lipid, TG, and CHO, chromium supplementation has a considerable impact on the obesity index, body fat, and blood lipid concentrations. In related research, Lewicki *et al.* [66] found that adding chromium ions (III) to a diet decreased blood glucose levels, LDL-c levels, cholesterol levels, and the risk of atherosclerosis and heart attacks.

Table 11. Liver cholesterol, total lipids, triglycerides (TGs), and glycogen of the obese investigational rat groups.

Parameters (mg/g)	Negative control	Positive control	Chromium	OPP	OPuP	OPP + chromium	OPuP + chromium	OPP+ OPuP+ chromium
CHO	4.01±0.39c	8.03±1.30a	6.11±0.96b	4.95±0.33c	4.88±0.55c	4.79±0.41c	4.99±0.53c	4.66±0.52c
T. lipids	34.77±3.14d	66.89±5.11a	55.77±5.91b	50.11±5.31b	46.50±4.91bc	45.61±4.77bc	44.75±4.35bc	40.77±4.3c
TGs	4.51±0.41a	2.01±0.25c	3.19±0.37b	4.45±0.45a	4.31±0.35a	4.20±0.42a	4.01±0.47a	4.91±0.44a
Glycogen	4.02±0.42a	2.11±0.27b	3.03±0.43a	3.49±0.26a	3.71±0.39a	3.61±0.37a	3.51±0.35a	3.66±0.34a

Each value represents the Mean \pm SD (n=6). Means under the same row with different superscript letters are significantly different at $p \leq 0.05$. OPP, orange peel powder; OPuP, orange pulp powder; CHO, cholesterol; T. lipids, total lipids; TGs, triglyceri

CONCLUSION

The current study has shown the effectiveness of specific food industry waste, namely peels and pulps, when combined with chromium to treat problems or abnormalities in obese rats. The liver functioning, hyperglycemia, and serum lipid profile are some of these consequences. Due to their high abundance of bioactive constituents, vitamins, and minerals, they exhibit a considerable antioxidant activity, which is responsible for all these benefits. These results lay the groundwork for the use of specific food industry by-products in the early treatment of obesity. Additionally, the current study suggested adding orange by-product powder and/or extracts to our everyday meals, beverages, dietary supplements, and pharmaceutical formulations.

CONFLICTS OF INTEREST

There is no conflict to declare.

REFERENCES

1. Beers, M.H. and Berkow, R. (2004): Nutritional Disorders: Obesity. Section 1, Chapter 5. In *The Merck Manual of Diagnosis and Therapy*. Whitehouse Station, NJ: Merck Research Laboratories.
2. Mehram, E.; Aboraya, A. and Elhassaneen, Y. (2021). Potential Effects of Food Processing Byproducts on Neurological and Immunological Disorders of Obese Rats. *Alexandria Science Exchange Journal*, 42, (2): 509-522. [DOI: 10.21608/asejaiqjsae.2021.178864].
3. Zahran, H., Gad, M. A., Fawzy, M., & Hassan, H. (2021). The role of different edible vegetable oils as hypolipidemic agents on experimental hyper-lipidemic rat model. *Egyptian Journal of Chemistry*, 64(11), 6231-6242.
4. Caterson, I.D. (2009). Medical Management of Obesity and its Complications. *Annals Academy of Medicine*, 38 (1):22-28.
5. Muñoz-Garach, A.; Isabel C. and Francisco J. (2016). Does Metabolically Healthy Obesity Exist? *Nutrients*, 8: 1-10.
6. OECD, Organization for Economic Co-operation and Development member countries (2017). Obesity Update 2017. www.oecd.org/health/obesity-update.htm.
7. WHO (2006). Report. Puska, P. Nishida, C. Porter, D. Obesity and over weight. Rome.
8. Callaway, L.; Prins, J.; Chang, A. and McIntyre, D. (2006): The prevalence and impact of overweight and obesity in an Australian obstetric population. *MJA*. 184(2):56-9.
9. IOM (Institute of Medicine).(2009): Weight gain during pregnancy; reexamining the guidelines. Washington DC: The National Academy Press (US).
10. Catalano, P.M. and Ehrenberg, H.M. (2006): The short- and long-term implications of maternal obesity on the mother and her offspring. *BJOG: An International Journal of Obstetrics & Gynaecology*.; 113(10):1126-33.
11. Birdsall, KM. Vyas, S. Khazaezadeh, N. Oteng-Ntim, E.(2009). Maternal obesity: a review of interventions. *Int J Clin Pract.*; 63(3):494-507.
12. WHO (1999). Definition, diagnosis and classification of diabetes mellitus and its complications. World Health Organization Department of Noncommunicable Disease Surveillance. (<http://whglidoc.who.int/hg/1999/WHO-NCD-NCS-99.2pdf>). 60 p. Retrieved on 7/6/2007.
13. Elabd, E. M. Y., Zahran, H. A., & Abdalla, A. M. (2017). A comparative study of the effects of three Moringa species on obesity-induced oxidative stress state in liver tissue. *Int J Pharma Bio Sci*, 8, 572-584.
14. Elhassaneen, Y.; Mekawy, S.; Khder, S. and Salman, M. (2019). Effect of Some Plant Parts Powder on Obesity Complications of Obese Rats. *Journal of Home Economics*, 29 (1): 83-106. DOI: 10.21608/mkas.2017.166177.
15. Elhassaneen, Y.; Ragab, S. and Essa E. (2020). Chemical and nutritional studies on extracts of food processing by-products and their effects on obesity complications in rats. *Journal of Home Economics*, 30 (2): 1-26. DOI: 10.21608/mkas.2020.156506.
16. Abd-Elwahed, H.M. (2022). Antioxidant activities and bioactive compounds content of some plant parts as fat burners: applications in obese rats. M.Sc. Thesis in Nutrition and Food Science, Faculty of Specific Education, Benha University, Benha, Egypt.
17. Velasco, R; Licciardello, C (2014). "A genealogy of the citrus family". *Nature Biotechnology*. 32 (7): 640–642.
18. Bampidis V., Robinson P. (2006): Citrus by-products as ruminant feeds: a review. *Animal Feed Science and Technology*, 128:175–217.
19. Mondello L.; Casilli A.; Tranchida P.; Dugo P. and Dugo G. (2005): Comprehensive two-dimensional GC for the analysis of Citrus essential oils. *Flavour and Fragrance Journal*, 20: 136-40.
20. M'hiri N.; Ioannou I.; Ghoul M. and Mihoubi N. (2015): Proximate chemical composition of orange peel and variation of phenols and antioxidant activity during convective air drying. *Journal of New Science*, 9: 881-90.
21. Ben Hsouna A., Hamdi N., Ben Halima N., Abdelkafi S. (2013): Characterization of essential oil from Citrus aurantium L. flowers: Antimicrobial and Antioxidant activities. *Journal of Oleo Science*, 62(10): 763-772.
22. Lv, Y.X.; Zhao, S.P.; Zhang, J.Y.; Zhang, H.; Xie, Z.H.; Cai, G.M. and Jiang, W.H. (2012). Effect of orange peel

- essential oil on oxidative stress in AOM animals. *Int J Biol Macromol.* 2012 May 1;50(4):1144-50.
23. AOAC. (1995). "Official Methods of the Association of Official analytical Chemists" 16 th Ed. Published by the Association of Official Analytical Chemists. Arlington, Virginia, VA.
 24. Singh, K.; Sundarro, K.; Tinkerame, J.; Kaluwin, C. and Matsuoka, T. (1991). Lipid content fatty acid and mineral composition of Mud Crabs (*Seylla serrata*) from Papua new Guinea. *Journal of Food Composition and Analysis*, 4 (3): 276 – 280.
 25. Moeslinger, T.; Brunner, M. and Spieckermann, G. (1994). Spectrophotometric determination of dehydroascorbic acid in biological samples. *Analytical Biochemistry*. 221: 290 - 296.
 26. Epler, K.S.; Zeigler, R.G. and Craft, N.E. (1993). Liquid chromatographic method for the determination of carotenoids, retinoids and tocopherols in human serum and in food. *J Chromatog.* 619:37–48.
 27. Hung, S.S.; Cho, Y.C. and Slinger, S.J. (1980): High performance liquid chromatographic determination of alpha-tocopherol in fish liver, *J. Assoc. Off. Anal. Chem.*, 63 (4): 889 - 893.
 28. Singleton, V. and Rossi, J. (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.*, 16: 144-158.
 29. Wolfe, K.; Wu, X. and Liu, R.H. (2003). Antioxidant activity of apple peels. *J Agric Food Chem.* 51:609–614.
 30. Lichtenthaler, H.K. (1987) Chlorophylls and Carotenoids, the Pigments of Photosynthetic Biomembranes. In: Douce, R. and Packer, L., Eds., *Methods in Enzymology*, Academic Press Inc., New York, 350-382.
 31. Zhishen, J.; Mengcheng, T. and Jianming, W. (1999). The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals, *Food Chemistry*, 64(4):555-559.
 32. Marco, G.J. (1968). A rapid method for evaluation of antioxidants. *J. Am. Oil Chem. Soc.*, 45: 594-598.
 33. Al-Saikhan, M. S.; Howard, L. R. and Miller, J. C., Jr. (1995). Antioxidant activity and total phenolics in different genotypes of potato (*Solanum tuberosum*, L.). *Journal of Food Science*. 60 (2): 341-343.
 34. AIN. American Institute of Nutrition. (1993): Purified diet for laboratory Rodent, Final report. *J. Nutrition*. 123:1939-1951.
 35. NRC, National Research Council (1996). *Guide for the Care and Use of Laboratory Animals* Washington: National Academy Press.
 36. Chapman, D.G.; Castilla, R. and Champbell, J.A. (1959): Evaluation of protein in food. I.A Method for the determination of protein efficiency ratio. *Can. J. Biochemistry and Physiology*, 37: 679-686.
 37. Drury, R .A. and Wallington, E.A. (1980): *Carleton's Histological Techniques*, 5th ed, Oxford University Press. London, New York Toronto: 1:653-661.
 38. Everds N. and Ramaiah L. (2017): *The Laboratory Rats*. In: Kurts D.M., Travlos G.S (eds): *The Clinical Chemistry of Laboratory Animals*. 3rd Ed. Milton, United Kingdom, Taylor & Francis Group: 33-78.
 39. Lee R., Nieman D. (1996): *Nutritional Assessment*. 2nd Ed. New York, USA, McGraw Hill:303-337.
 40. Castelli, T. and Levitar, Y. (1977): *Atherogenic*, index *Curr Presc*: p39
 41. Snedecor, G. W. and Cochran, W. G. (1967): *Statistical Methods*" 6th Ed. Iowa State, USA.
 42. M'hiri N., Ioannou I., Ghoul M., Mihoubi N. (2014). *Extraction Methods of Citrus Peel Phenolic Compounds*. *Food Reviews International*, 30 (4): 265–90.
 43. Forrest, H.N. (2000). Importance of making dietary recommendations for elements designated as nutritionally beneficial, pharmacologically beneficial, or conditionally essential. In: *Special Issue:Trace Elements in Human Health and Disease: An Update*, *The Journal of Trace Elements in Experimental Medicine*, 13 (1): 113-129.
 44. Osredkar, J. and Sustar, N. (2011). Copper and zinc, biological role and significance of copper/zinc imbalance. *J. Clinic Toxicol. S.* S3:001.
 45. Nazanin, A.; Richard, H. and Roya, K. (2014). Review on iron and its importance for human health. *J. Res. Med. Sci.* 19(2):164-174.
 46. Elhassaneen, Y.; El-Waseef, S.; Fathy, N. and Sayed Ahmed, S. (2016): Bioactive compounds and antioxidant potential of food industry by-products in Egypt. *American Journal of Food and Nutrition*, 4 (1): 1-7. DOI:10.12691/ajfn-4-1-1.
 47. Velioglu, Y.S.; Mazza, G.; Gao, L. and Oomah, B.D. (1998). Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *J. Agric. Food Chem.*, 46 (10): 4113-4117.
 48. Hegazy, W. H. (2009). Antioxidant activity of pomegranate (*punica granatum*) fruits and its relationship with phenolic composition and processing technology. M.Sc. Thesis in Nutrition and Food Science, Faculty of Home Economics, Minoufiya University, Shebin El-Kom, Egypt.
 49. El-Mokadem, K. (2010). The effect of technological treatments on phytochemical properties of some foods ". M.Sc. Thesis in Nutrition and Food Science, Faculty of Home Economics, Minoufiya University, Egypt.
 50. Sayed Ahmed, S. (2016). Nutritional and technological studies on the effect of phytochemicals on obesity injuries and their related diseases by using experimental animals ". Ph.D. Thesis in Nutrition and Food Science, Faculty of Specific Education, Port Saied University, Port Saied, Egypt

51. Halliwell, B. (1991): Reactive oxygen species in living systems: Source, biochemistry and role in human disease. *American Journal of Medicine*, 91: 14s-21s.
52. Chaitanya, K.V.; Pathan, A.A.K.; Mazumdar, S.S.; Charavarthi, G.P.; Parine, N. and Bobbarala, V. (2010): Role of oxidative stress in human health: An overview. *Journal of Pharmacy Research*. 3: 1330-1333.
53. Elmaadawy A., Arafa R. and Elhassaneen Y. (2016). Oxidative Stress and antioxidant defense systems status in obese rats feeding some selected food processing by-products applied in bread. *Journal of Home Economics*, 26 (1): 1-38.
54. Alqallaf, J. H. (2021). "Effect of feeding fortified by some Brassicaceae family vegetables on biological changes of obesity rats induced". Ph.D. Thesis in Nutrition and Food Science, Faculty of Specific Education, Benha University, Benha, Egypt
55. Rodriguez, D. S.; Hadley, M. and Holm, E. T. (1994): Potato peel waste: Stability and antioxidant activity of a freeze-dried extract. *Journal of Food Science*. 59(5): 1031-1033.
56. Beattie, J.; Crozier, A. and Duthie, G. (2005). Potential Health Benefits of berries. *Current Nutrition and Food Science*. 1: 71-86.
57. Kim B., Lindemann M., Cromwell G. (2010): Effects of dietary chromium (III) picolinate on growth performance, respiratory rate, plasma variables, and carcass traits of pigs fed high-fat diets. *Biological Trace Element Research*, 133(2):181-196.
58. Liu Y.; Heying E. and Tanumihardjo S. (2012): History, global distribution, and nutritional importance of citrus fruits. *Comprehensive Reviews in Food Science Food Safety*, 11:530-45.
59. Osfor M., Hegazy A., El-Moaty M., Elmadbouly M., Afify R., Elbahnasawy A. (2013): Hypo-cholesterolemic and hypoglycemic effects of orange albedo powder (citrus aurantiuml) on male albino rats. *International Journal of Nutrition and Food Sciences*, 2(2): 70-76.
60. Avenell, A.; Broom, J.; Brown, T.J.; Poobalan, A.; Aucott, L. and Stearns, S.C. (2004): Systematic review of the long-term effects and economic consequences of treatments for obesity and implications for health improvement. *Health Technology Assess*, 8 (21): 37-49.
61. Bales, C.W. and Buhr, G. (2008): Is obesity bad for older persons? A systematic review of the pros and cons of weight reduction in later life. *J Am Med Dir Assoc*; 9(5):302-12.
62. Williamson, D.A.; Rejeski, J.; Lang, W.; Van Dorsten, B.; Fabricatore, A.N. and Toledo, K. (2009): Impact of a weight management program on health-related quality of life in overweight adults with type 2 diabetes. *Arch Intern Med.*, 169(2):163-171.
63. Racek J.; Trefil L.; Rajdl D.; Mudrova V.; Hunter D. and Senft, V. (2006): Influence of chromium-enriched yeast on blood glucose and insulin variables, blood lipids, and markers of oxidative stress in subjects with type 2 diabetes mellitus. *Biological Trace Element Research*, 109(3): 215-230.
64. Sahin K., Onderci M., Tuzcu M., Ustundag B., Cikim G., Ozercan I., Sriramaju V., Juturu V., Komorowski J. (2007): Effect of chromium on carbohydrate and lipid metabolism in a rat model of type 2 diabetes mellitus: the fat-fed, streptozotocin-treated rat. *Metabolism Clinical and Experimental*, 56(9):1233-1240.
65. Tiwari A.; Saha S.; Yadav V.; Upadhyay K.; Katiyar D., and Mishra T. (2017): Extraction and characterization of pectin from orange peels. *International Journal of biotechnology and Biochemistry*, 13(1):39-47.
66. Lewicki S., Zdanowski R., Krzyzowska M., Lewicka A., Dębski B., Niemcewicz M., Goniewicz M. (2014): The role of Chromium III in the organism and its possible use in diabetes and obesity treatment. *Annals of Agricultural and Environmental Medicine*, 21(2): 331-335.